arXiv:1208.1881v1 [math.DS] 9 Aug 2012

POLYNOMIAL SIEGEL DISKS ARE TYPICALLY JORDAN DOMAINS

GAOFEI ZHANG

Abstract. We prove that for typical rotation numbers polynomial Siegel disks are Jordan domains with boundaries containing at least one critical point.

1. Introduction

Let f be a holomorphic germ which has an indifferent fixed point. Suppose f is linearizable at p. The maximal linearizable domain D containing p is called the Siegel disk of f centered at p. The topology of the boundaries of Siegel disks is one of the important topics in complex dynamics. For general holomorphic germs, the boundary of the Siegel disk can be very complicated. Recently Chéritat constructed a holomorphic germ with a Siegel disk compactly contained in the germ such that the boundary of the Siegel disk is non-locally connected [2]. For rational maps, however, it has been believed that the Siegel disk is always a Jordan curve. This was conjectured by Douady and Sullivan in early 1980's and is one of the dominant open problems in this area. The conjecture has been settled in the case that the rotation number of the Siegel disk is of bounded type. Here we say an irrational number $0 < \theta < 1$ is of *bounded type* if all the coefficients of its continued fraction have a finite upper bound.

Theorem 1 (Douady-Herman [7], Zakeri [22], Shishikura [18], Zhang [23])**.** *All bounded type Siegel disks of rational maps are quasi-disks with at least one critical point on their boundaries.*

In contrast with the bounded type condition imposed on the rotation numbers, Ino and Shishikura recently showed that the conjecture holds for quadratic Siegel disks provided that the coefficients of the continued fractions are all large enough.

Theorem 2 (Inou-Shishikura [10])**.** *There is an $N > 1$ such that for any Brjuno number θ, if all the coefficients of the continued fraction of θ are larger than N, then the boundary of the Siegel disk of $P_\theta(z) = e^{2\pi i\theta}z + z^2$ is a Jordan curve.*

Note that the sets of both the bounded type and Ino-Shishikura type irrational numbers have zero Lebesgue measure.

In 2002 Petersen and Zakeri proved that for typical rotation numbers, a quadratic Siegel disk is a Jordan domain with a critical point on its boundary. To state their theorem more precisely, let us introduce another type of irrational numbers. Let $C > 0$. Let Θ_C denote the set of all irrational numbers $0 < \theta < 1$ such that

$$\log a_n \leq C\sqrt{n} \tag{1}$$

2000 *Mathematics Subject Classification.* 58F23, 37F10, 37F45, 32H50, 30D05.

holds for all $n \geq 1$ where $[a_1, a_2, \cdots]$ is the continued fraction of θ. We say $0 < \theta < 1$ is of *David type* if $\theta \in \Theta_C$ for some $C > 0$. It is known that the set of David type irrational numbers in $[0, 1]$ has full Lebesgue measure [11].

Theorem 3 (Petersen-Zakeri [17])**.** *Let $0 < \theta < 1$ be a David type irrational number. Then the Siegel disk of $P_\theta(z) = e^{2\pi i\theta}z + z^2$ is a Jordan domain whose boundary contains the unique finite critical point of P_θ.*

The main purpose of this paper is to generalize the above result to polynomial maps of all degrees.

Main Theorem. *All David type Siegel disks of polynomial maps are Jordan domains with boundaries containing at least one of the critical points.*

One of the fundamental tools in our proof is trans-qc surgery. This surgery technique was pioneered by Haissinsky, who used it to transform an attracting basin into a parabolic basin, and then introduced to the study of Siegel disks by Petersen and Zakeri in [17]. Compared with qc surgery, the main difficulty in performing trans-qc surgery is to verify the integrability of certain degenerate Beltrami differentials. This often requires some delicate area estimates. In [17] the authors there used Petersen puzzles to obtain the desired estimate for the Douady-Ghys premodel. This idea was then adapted to some slightly more general situations in [25]. Up to now it is not known if the invariant Beltrami differential is always integrable for an arbitrarily given premodel. This is the essential challenge in generalizing Petersen-Zakeri's theorem to polynomial maps of all degrees.

The idea of our proof can be sketched as follows. Suppose $C > 0$ is a fixed constant and D is a David type Siegel disk of an arbitrary polynomial map with rotation number $\theta \in \Theta_C$. By perturbing θ we get a sequence of bounded type Siegel disks D_N with rotation numbers $\theta_N \in \Theta_C$ such that $\theta_N \to \theta$. By Shishikura's theorem, each ∂D_N is a quasi-circle passing through at least one of the critical points. We shall see if θ is not of bounded type, the qc constants of $\partial D_N, N \geq 1$, are not bounded. This means that the oscillations of these quasi-circles can not be uniformly controlled with respect to the qc constants. Thus nothing can be obtained if we let N go to ∞ at this point. The key of our proof is to find an appropriate way to measure the oscillations of these quasi-circles so that in this way, the oscillations can be uniformly controlled. To do this, we will introduce a sequence of oscillation functions. We prove that these oscillation functions are uniformly bounded for bounded type Siegel disks of a class of special polynomial maps with the rotation numbers belonging to Θ_C. This is the place where we use trans-qc surgery. We then show that for a bounded type Siegel disk of an arbitrary polynomial map with the rotation number belonging to Θ_C, the oscillation functions can be controlled, in certain sense, by those for the special ones. From this we derive that the oscillations of the sequence of quasi-circles are uniformly bounded. By passing to a subsequence if necessary, we may assume that the sequence of quasi-circles converge to some Jordan curve which passes through at least one of the critical points. This Jordan curve must be the boundary of the David type Siegel disk. This proves the Main Theorem.

The following is the organization of the paper.

In §2 we present a detailed outline of the proof. We first formulate a reduced version of the Main Theorem by introducing oscillation functions. We then state four key lemmas.

The proofs of these four key lemmas consist of the core part of the paper. Finally we prove the Reduced Main Theorem by assuming the four key lemmas.

In §3 we prove that the Reduced Main Theorem implies the Main Theorem.

In §4 we prove the first key lemma. This lemma says that the oscillation of the boundaries of bounded type Siegel disks for a class of special polynomial maps, with rotation numbers belonging to Θ_C, are uniformly bounded. The proof uses essentially the idea of trans-qc surgery.

In §5 we prove the third and fourth key lemmas. These two lemmas are used to construct a chain of slices in the parameter space. Each of these slices is an algebraic Riemann surface determined by a finite system of polynomial equations. We use this chain of Riemann surfaces to establish a connection between an arbitrary Siegel polynomial map to those special ones.

In §6 we establish a topological characterization of a class of polynomial maps with bounded type Siegel disks. It is an extension of Thurston's characterization for post-critically finite rational maps. As a direct consequence, we prove the Key-Lemma 2. We use this lemma to perturb certain Siegel polynomial map so that the resulted one can be embedded into an appropriate slice in the parameter space.

In §7, the Appendix of the paper, we present a list of basic properties about bounded type Siegel disks of polynomial maps. Among these the most important one is Shishikura's theorem that all bounded type Siegel disks of polynomial maps are quasi-disks with qc constants depending only on the degree and the rotation number. From this we easily derive that when the rotation number is fixed, the boundary of bounded type Siegel disks moves continuously. This property plays an important role in our proof.

Acknowledgement. I would like to express my deep thanks to Prof. Carsten Lunde Petersen who spent many hours discussing with me on an early version of the manuscript during his visit of Nanjing in March, 2012.

2. Outline of the proof

Fix an integer $d \geq 2$ and a David type irrational number $0 < \theta < 1$ throughout the paper. Let $[a_1, \cdots, a_n, \cdots]$ be the continued fraction of θ. By definition we have $C > 0$ such that

$$\log a_n \leq C\sqrt{n}$$

for all $n \geq 1$. Let

$$P(z) = e^{2\pi i\theta}z + \alpha_2 z^2 + \cdots + \alpha_d z^d$$

with $\alpha_d \neq 0$. We want to show that the Siegel disk of P centered at the origin is a Jordan domain with at least one critical point on its boundary.

For the above $C > 0$, recall that Θ_C denotes the set of all the irrational numbers $0 < \theta < 1$ such that $\log a_n \leq C\sqrt{n}$ for all $n \geq 1$. Let

$$\Theta_C^b \subset \Theta_C$$

denote the subset consisting of all the bounded type irrational numbers in Θ_C. Let $\theta_N \in \Theta_C^b$, $N \geq 1$, be a sequence such that $\theta_N \to \theta$ as $N \to \infty$. Such sequence can be constructed in many ways. For instance, we can define

$$\theta_N = [a_1, \cdots, a_N, 1, 1, 1, \cdots].$$

For each $N \geq 1$, let

$$P_N(z) = e^{2\pi i\theta_N} z + \alpha_2 z^2 + \cdots + \alpha_d z^d.$$

Then P_N converges to P uniformly in any compact set of the complex plane. It follows that the critical sets of all P_N are contained in a neighborhood of that of P, and therefore contained in a compact set of the plane. Let D_N denote the Siegel disk of P_N centered at the origin. Since θ_N is of bounded type, by Shishikura's theorem ([18], see also [23]), there is a critical point c_N of P_N and a $K_N > 1$ depending only on

$$\sup_{1 \leq k \leq N} \{a_k\},$$

such that ∂D_N is a K_N-quasi-circle and passes through c_N. By taking a subsequence, we may assume that c_N converges to some critical point c of P. Note that $K_N \to \infty$ if $\sup_{k\geq 1}\{a_k\} = \infty$. Because otherwise, by taking a subsequence, ∂D_N would converge to a quasi-circle passing through c. This quasi-circle must be the boundary of the Siegel disk of P centered at the origin. But by a result of Petersen [16], the rotation number of such Siegel disk must be of bounded type. This is a contradiction.

Let

$$Q = c^{-1}P(cz) \text{ and } Q_N(z) = c_N^{-1}P(c_N z).$$

Then 1 is a critical point of both Q and Q_N. Let us still use D_N to denote the Siegel disk of Q_N centered at the origin. It follows that $1 \in \partial D_N$ for all $N \geq 1$.

The main task of our proof is to show that the sequence of curves ∂D_N converge to a Jordan curve passing through the critical point 1. Since the quasiconformal constants K_N is unbounded, we need to find an appropriate way to measure the oscillation of these curves such that in this way, the oscillation of the curves can be uniformly controlled. This is the key of the work. More precisely,

Reduced Main Theorem. *There exist a pair of positive functions* $\eta, \lambda : (0, 2] \to \mathbb{R}^+$ *satisfying*

$$\lim_{\delta \to 0_+} \lambda(\delta) = \lim_{\delta \to 0_+} \eta(\delta) = 0$$

such that for any $N \geq 1$*, any pair of integers* $k > m \geq 0$*, and any pair of positive numbers* $0 < \delta' < \delta$*, if*

$$\delta' < |e^{ik\theta_N} - e^{im\theta_N}| < \delta,$$

then

$$\eta(\delta') \leq |Q_N^k(1) - Q_N^m(1)| \leq \lambda(\delta). \tag{2}$$

Proposition 2.1. The Reduced Main Theorem implies the Main Theorem.

The proof of Proposition 2.1 will be given in §3. The main task of the paper is to prove the Reduced Main Theorem. The proof is based on four key lemmas.

Definition 2.1. Let $0 < \alpha < 1$ be a bounded type irrational number. Let

$$\Pi_{\text{geom}}^{\alpha,d}$$

denote the class of all polynomial maps f of degree d such that

1. f has a Siegel disk centered at the origin with rotation number α,
2. the point 1 is a critical point of f,
3. all the finite critical points of f are contained in the boundary of the Siegel disk.

We will give a topological characterization of a more general class of Siegel polynomial maps (cf. Theorem 2.1). We will see that each $f \in \Pi_{\rm geom}^{\alpha,d}$ is uniquely determined by the $d-2$ angles between 1 and all the other $d-2$ critical points.

The first key lemma states that the Reduced Main Theorem holds for these special Siegel polynomial maps. Recall Θ_C^b is the subset of Θ_C consisting of all bounded type irrational numbers in Θ_C.

Key-Lemma 1. *There exist a pair of positive functions* $\eta_1, \lambda_1 : (0,2] \to \mathbb{R}^+$ *satisfying*

$$\lim_{\delta\to 0_+} \lambda_1(\delta) = \lim_{\delta\to 0_+} \eta_1(\delta) = 0,$$

such that for any

$$f \in \bigcup_{\alpha\in\Theta_C^b} \Pi_{\rm geom}^{\alpha,d},$$

any pair of integers $k > m \geq 0$ *and any pair of positive numbers* $0 < \delta' \leq \delta$ *satisfying* $\delta' \leq |e^{ik\theta} - e^{im\theta}| \leq \delta$, *the inequality*

$$\eta_1(\delta') \leq |f^k(1) - f^m(1)| \leq \lambda_1(\delta)$$

holds.

The fundamental tool in the proof of Key-Lemma 1 is trans-qc surgery. The proof is based on Lemmas 2.1- 2.3. Before we state the three lemmas, let us introduce some notations first. For any $f \in \bigcup_{\alpha\in\Theta_C^b} \Pi_{\rm geom}^{\alpha,d}$, we will show that there is a Blaschke product B_f of degree $(2d-1)$ which models f. In particular, all the critical points of B_f, except 0 and ∞, are contained in $\mathbb{T}$. Let $h_f : \mathbb{T} \to \mathbb{T}$ be the circle homeomorphism such that

$$h_f \circ (B_f|\mathbb{T}) \circ h_f^{-1} = R_\alpha.$$

Note that α is of bounded type and $h_f : \mathbb{T} \to \mathbb{T}$ is a quasisymmetric circle homeomorphism. Because the qc constants can not be uniformly controlled, instead of making a usual quasiconformal extension of h_f and then performing a qc surgery, we will construct a David extension $H_f : \Delta \to \Delta$ of h_f by adapting the idea in [17] and then perform a trans-qc surgery. Since α is of bounded type, the map H_f obtained in this way is necessarily quasiconformal. The key point here is that we regard H_f as a David homeomorphism when we measure its distortion.

Lemma 2.1. *There exist* $M, \beta > 0$ *and* $0 < \epsilon_0 < 1$ *depending only on* C *and* d *such that for any* $f \in \bigcup_{\alpha\in\Theta_C^b} \Pi_{\rm geom}^{\alpha,d}$, *the conjugation map* $h_f : \mathbb{T} \to \mathbb{T}$ *has a David extension* $H_f : \Delta \to \Delta$ *satisfying the following. For any* $0 < \epsilon < \epsilon_0$, *we have*

$$area\{z \in \Delta \mid |\mu_{H_f}(z)| > 1 - \epsilon\} < Me^{-\frac{\beta}{\epsilon}}.$$

The key idea in the proof of Lemma 2.1 is that h_f has a uniform saddle-node geometry. One of the most essential tools in our proof is Herman's uniform estimate on the distortion of cross-ratios for the compact family

$$\{B_f \mid f \in \bigcup_{\alpha\in\Theta_C^b} \Pi_{\rm geom}^{\alpha,d}\}. \tag{3}$$

We would like to point out that the compactness of the family (3) plays an essential role in the proof. The corresponding family in constructing Siegel disks of rational maps is not compact, even for quadratic rational maps. In this case Herman's uniform estimate does not hold and the conjugation map h_f does not have uniform saddle-node geometry. We believe Lemma 2.1 still holds though we do not know how to prove it. This is the most essential difficulty to overcome if one wants to use the idea of this paper to generalize the Main Theorem to all rational maps.

Now let $H_f : \Delta \to \Delta$ be the David homeomorphism in Lemma 2.1. Define

$$\widehat{B}_f(z) = \begin{cases} B_f(z) & \text{for } z \in \widehat{\mathbb{C}} \setminus \Delta, \\ H_f^{-1} \circ R_\alpha \circ H_f(z) & \text{for } z \in \Delta. \end{cases}$$

Let μ_f denote the Beltrami differential on the whole plane which is obtained by pulling back μ_{H_f} through the iteration of $\widehat{B}_f$.

Lemma 2.2. *There exist $\tilde{M}, \tilde{\beta} > 0$ and $0 < \tilde{\epsilon}_0 < 1$ depending only on C and d such that for any $f \in \bigcup_{\alpha \in \Theta_C^b} \Pi_{\text{geom}}^{\beta,d}$, we have*

$$area\{z \in \mathbb{C} \mid |\mu_f(z)| > 1 - \epsilon\} < \tilde{M} e^{-\frac{\tilde{\alpha}}{\epsilon}}$$

for any $0 < \epsilon < \tilde{\epsilon}_0$.

Lemma 2.2 asserts the uniform integrability of the invariant Beltrami differentials for all those premodels whose finite critical points are all contained in the unit circle. As we mentioned before the main difficulty in performing a trans-qc surgery is to verify the integrability of certain degenerate Beltrami differential. The key idea used in the proof of Lemma 2.2 is a method developed in [25] which allows us to obtain the desired area estimate.

Lemma 2.3 (Tukia, [19])**.** *Let $\mathcal{F}$ denote the class of all $(\tilde{M}, \tilde{\beta}, \tilde{\epsilon}_0)$-David homeomorphisms of the plane to itself which fix 0 and 1. Then there exist three constants $\hat{M}, \hat{\beta} > 0$ and $0 < \hat{\epsilon}_0 < 1$ depending only on $\tilde{M}, \tilde{\beta}$ and $\tilde{\epsilon}_0$ such that any sequence in $\mathcal{F}$ has a subsequence which converges uniformly to a $(\hat{M}, \hat{\alpha}, \hat{\epsilon}_0)$-David homeomorphism of the plane which fixes 0 and 1.*

The Key-Lemma 1 is a direct consequence of Lemmas 2.1 - 2.3 (see §4).

We would like to point out that in the case of cubic polynomial maps, the Reduced Main Theorem follows from Key-Lemma 1. The following is the detailed argument.

Let $\alpha \in \Theta_C^b$ and let $\mathcal{P}_\alpha^3$ denote the space of all cubic polynomials which have a Siegel disk at the origin with rotation number α and a critical point at 1. Let c denote the other critical point. Then $\mathcal{P}_\alpha^3$ is parameterized by c and under this parametrization, $\mathcal{P}_\alpha^3$ is homeomorphic to the punctured plane $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$. For each $c \in \mathbb{C}^*$, let f_c denote the corresponding cubic polynomial and D_c denote the Siegel disk of f_c centered at the origin. For any two integers $k > m \geq 0$, define

$$\sigma_{k,m}(c) = f_c^k(1) - f_c^m(1).$$

Since f_c depends holomorphically on c when c varies in $\mathbb{C}^*$, $\sigma_{k,m}$ is a holomorphic function in $\mathbb{C}^*$. By a result of Zakeri (cf. §14 of [22]), there is a Jordan curve $\Gamma \subset \mathbb{C}^*$ such that

1. Γ separates 0 and ∞, passes through 1 and is invariant under $c \to 1/c$,
2. for all c belonging to the inside of Γ and not equal to 0, ∂D_c passes through the critical point c only; for all c belonging to the outside of Γ and not equal to ∞,

∂D_c passes through the critical point 1 only; for all c belonging to Γ, ∂D_c passes through both 1 and c.

For the α given and $d = 3$, let λ_1 and η_1 be the two positive functions in the Key-Lemma 1. Note that Γ corresponds to the class $\prod_{\text{geom}}^{\alpha,3}$ by the second assertion above. So for any pair of integers $k > m \geq 0$ and any pair of positive numbers $0 < \delta' \leq \delta$ satisfying $\delta' \leq |e^{ik\theta} - e^{im\theta}| \leq \delta$, we have

$$\eta_1(\delta') \leq |\sigma_{k,m}(c)| \leq \lambda_1(\delta) \quad \text{for all } \ c \in \Gamma. \tag{4}$$

Since 1 belongs to ∂D_c for c belonging to the outside of Γ, $\sigma_{k,m}$ does not vanish in the outside of Γ. Noting that as $c \to \infty$, f_c converges uniformly to a quadratic polynomial, it follows that $\sigma_{k,m}$ has a removable singularity at infinity. This implies $\sigma_{k,m}$ is a holomorphic function in the outside of Γ and doers not vanish. So both the maximal and minimal principle applies. It follows that (4) holds for all c belonging to the outside of Γ. The Reduced Main Theorem for cubic polynomials follows by taking $\lambda = \lambda_1$ and $\eta = \eta_1$.

The argument above, however, does not work for polynomial maps of degree $d \geq 4$. The reason is as follows. For each $\alpha \in \Theta_C$, a Siegel polynomial map in $\Pi_{\text{geom}}^{\alpha,d}$ is uniquely determined by the $d-2$ angles between 1 and all the other $d-2$ finite critical points. More precisely, there is a natural homeomorphism between $\Pi_{\text{geom}}^{\alpha,d}$ and the set

$$\underbrace{S^1 \times \cdots \times S^1}_{(d-2)\ \text{copies}} / G_{d-2}$$

where G_{d-2} is the permutation group of order $d-2$. This implies that the set $\Pi_{\text{geom}}^{\alpha,d}$ has real dimension $d-2$. Thus for $d \geq 4$, the set $\Pi_{\text{geom}}^{\alpha,d}$ can not bound a domain in the parameter space

$$\underbrace{\mathbb{C}^* \times \cdots \times \mathbb{C}^*}_{(d-2)\ \text{copies}},$$

and we can not use maximal and minimal principles directly. To solve this problem, we will introduce certain slices in the parameter space. Each slice is an algebraic Riemann surface determined by a system of polynomial equations. We will apply the maximal and minimal principles successively on a chain of such slices, and finally prove that the oscillation of the boundary a Siegel disk for an arbitrary polynomial map, in certain sense, can be controlled by the oscillation of the boundaries of Siegel disks for those Siegel polynomial maps in the class $\Pi_{\text{geom}}^{\alpha,d}$. Since we have proved that the later can be uniformly controlled, the Main Theorem follows. The construction of these Riemann surfaces relies on the other three key lemmas. Before we state them, let us first introduce some notations.

Definition 2.2. Let $0 < \alpha < 1$ be an irrational number of bounded type and $d \geq 2$ be an integer. We use

$$\Sigma_{\text{geom}}^{\alpha,d}$$

to denote the class of all degree d polynomial maps with a Siegel disk D of rotation number α such that each critical point either is attracted to some periodic attracting cycle, or belongs to the boundary of the Siegel disk.

It is clear that

$$\Pi_{\text{geom}}^{\alpha,d} \subset \Sigma_{\text{geom}}^{\alpha,d}.$$

In §6 we will establish a topological characterization of all the polynomial maps in $\Sigma_{\text{geom}}^{\alpha,d}$. Suppose $g \in \Sigma_{\text{geom}}^{\alpha,d}$ such that the boundary of the Siegel disk contains more than one critical point. The Key-Lemma 2 states that one can always perturb g in the class $\Sigma_{\text{geom}}^{\alpha,d}$ so that after the perturbation all critical points in the boundary of the Siegel disk satisfy an orbit relation.

Key-Lemma 2. *Let $g \in \Sigma_{\text{geom}}^{\alpha,d}$. Suppose g has two or more distinct critical points on the boundary of the Siegel disk, say $c_1, \cdots, c_m$, where $m \ge 2$. Then for any $\epsilon > 0$, there is a $\tilde{g} \in \Sigma_{\text{geom}}^{\alpha,d}$ such that*

1. *$\|g-\tilde{g}\| < \epsilon$ where $\|g-\tilde{g}\|$ denotes the maximal absolute value of all the coefficients of $g - \tilde{g}$,*
2. *$\tilde{g}$ has exactly m distinct critical points on the boundary of the Siegel disk, say $\tilde{c}_i$, $1 \le i \le m$, such that $|c_i - \tilde{c}_i| < \epsilon$ for all $1 \le i \le m$,*
3. *there are integers k_i, $2 \le i \le m$, such that $\tilde{g}^{k_i}(\tilde{c}_1) = \tilde{c}_i$ for $2 \le i \le m$.*

The proof of Key-Lemma 2 is based on Theorem 2.1 which gives a topological characterization of all the maps in $\Sigma_{\text{geom}}^{\alpha,d}$. The theorem is an extension of Thurston's characterization theorem for post-critically finite rational maps. Before we state the theorem, let us introduce some terminologies first. A topological polynomial f of degree d is a branched covering map of the sphere to itself such that $f^{-1}(\infty) = \{\infty\}$. We say $\mathcal{O} = \{x_1, \cdots, x_p\}$ is a holomorphic attracting cycle of f if $\mathcal{O}$ is a periodic cycle of f of period p, and moreover, there is an open neighborhood U of $\mathcal{O}$ on which f is holomorphic and $|Df^p(x_1)| < 1$.

Definition 2.3. Let $0 < \alpha < 1$ be a bounded type irrational number. Let $\Sigma_{\text{top}}^{\alpha,d}$ denote the class of all topological polynomials of degree d such that

1. f has at least one critical point in $\mathbb{T}$,
2. $f|\Delta$ is the rigid rotation given by $z \to e^{2\pi i\alpha}z$,
3. any critical point of f either is attracted to some holomorphic attracting cycle of f, or belongs to $\mathbb{T}$.

Let P_f denote the closure of the union of all critical orbits of f. We say a map $f \in \Sigma_{\text{top}}^{\alpha,d}$ is CLH-equivalent to a map $g \in \Sigma_{\text{geom}}^{\alpha,d}$ if there exist two homeomorphisms ϕ and ψ of the plane to itself such that

1. $\phi|\Delta = \psi|\Delta$ is holomorphic,
2. for each holomorphic attracting cycle $\mathcal{O}$, there is an open neighborhood U of $\mathcal{O}$ such that $\phi|U = \psi|U$,
3. ϕ is isotopic to ψ rel $P_f \cup \cup_i U_i$ where $U_i s$ are open neighborhoods of all holomorphic attracting cycles,
4. $\phi \circ f = g \circ \psi$.

Theorem 2.1. *A map $f \in \Sigma_{\text{top}}^{\alpha,d}$ is CLH-equivalent to a map $g \in \Sigma_{\text{geom}}^{\alpha,d}$ if and only if f has no Thurston obstructions in the exterior of Δ. Such g if exists, must be unique up to a linear conjugation.*

For the definition of Thurston obstructions, the proof of Theorem 2.1, and how to derive the Key-lemma 2 from Theorem 2.1, see §6. Let us briefly sketch the idea of the proof of the Key-Lemma 2. Starting from g we get its topological model $f \in \Sigma_{\text{top}}^{\alpha,d}$.

Then f has no Thurston obstruction in the exterior of Δ. By perturbing f in a small neighborhood of $\mathbb{T}$ we get a sequence f_n in $\Sigma_{\rm top}^{\alpha,d}$ such that all the critical points of f_n which belong to the boundary of the Siegel disk satisfy some prescribed orbit relation. It is clear that f_n has no Thurston obstruction in the exterior of Δ. By Theorem 2.1, there is a sequence $g_n \in \Sigma_{\rm geom}^{\alpha,d}$ such that f_n is Thurston equivalent to g_n. In particular, all critical points of g_n which belong to the boundary of the Siegel disk satisfy the desired orbit relations. We then show that, by appropriate linear conjugations if necessary, the sequence g_n is pre-compact and the limit polynomial map is equal to g. This implies Key-Lemma 2.

Before we state the other two key lemmas, let us present some preliminaries first. Let $0 < \alpha < 1$ be a bounded type irrational number. Let $\mathcal{P}_\alpha^d$ denote the class of all the polynomial maps f such that

$$f(z) = e^{2\pi i\alpha} z + \alpha_2 z + \cdots + \alpha_d z^d$$

with $\alpha_d \neq 0$ and $f'(1) = 0$. Since f has $d-1$ critical points (counting by multiplicities) and at least one of them is contained in the boundary of the Siegel disk centered at the origin, there are at most $d-2$ attracting periodic cycles. Note that such f is uniquely determined by the set of its critical points. More precisely, for each $(d-1)$-tuple

$$X = (c_1, \cdots, c_{d-1}),\ \ c_i \in \mathbb{C}^* \text{ for } 1 \le i \le d-2, \text{ and } c_{d-1} = 1,$$

there is a unique $f \in \mathcal{P}_\alpha^d$ such that X is the critical set of f. By a simple calculation, we have

$$f(z) = \sum_{i=1}^{d} a_i z^i$$

with

$$a_i = e^{2\pi i\alpha} \cdot \left(\frac{(-1)^{i-1}}{i} \right) \cdot \frac{Q_{d-i}(c_1, \cdots, c_{d-1})}{c_1 \cdots c_{d-1}}$$

where Q_{d-i} is the degree-$(d-i)$ elementary polynomials of $c_1, \cdots, c_{d-1}$. Let us denote such f by

$$f_{c_1, \cdots, c_{d-2}, 1} \text{ or } f_X.$$

Let us recall some basic notions of algebraic functions. For more knowledge in this aspect, the reader may refer to [1]. Suppose $P(w,z)$ is an irreducible polynomial. Suppose P has degree m with respect to w. That is,

$$P(w,z) = b_0(z) w^m + b_1(z) w^{m-1} + \cdots + b_m(z)$$

where $b_i(z)$ are polynomials of z and $b_0(z)$ is not identically zero. Let $R(z)$ be the resultant of $P(w,z)$ and $P_w(w,z)$. Let

$$Z = \{z \in \mathbb{C} \mid R(z) = 0 \text{ or } b_0(z) = 0\}.$$

Then for any $z \in \mathbb{C} \setminus Z$, there are exactly m distinct w such that $P(w,z) = 0$. Thus the equation

$$P(w,z) = 0$$

determines a multi-valued analytic function $w = w(z)$ in the sense $P(w(z), z) = 0$. For each $z_0 \in Z$, there are two cases.

In the first case, $R(z_0) = 0$. Then there is a $w_0 \in \mathbb{C}$ such that $P(w_0, z_0) = 0$ and $P_w(w_0, z_0) = 0$. In this case, there is an integer $p \geq 2$ such that in a small neighborhood of (w_0, z_0),

$$w(z) = \sum_{k=0}^{\infty} a_k (z - z_0)^{k/p}.$$

In the second case, $b_0(z) = 0$. Then there are $l < m$ finite complex numbers w, counting by multiplicity, such that $P(w, z_0) = 0$. In this case, for z near z_0 and w near infinity,

$$w(z) = \sum_{k=-n}^{\infty} a_k (z - z_0)^{k/p}$$

where $p = m - l$ and $n \geq 1$ is some integer.

The set

$$\{(w, z) \in \mathbb{C}^2 \mid P(w, z) = 0, z \notin Z\}$$

consist of a Riemann surface embedded in $\mathbb{C}^2$. Note that for each $z_0 \in Z$, there may exist $w_0 \in \mathbb{C}$ such that $P(w_0, z_0) = 0$ and in a small neighborhood of (w_0, z_0), $w(z)$ is an analytic function of z. After we add all such points to the above set, the resulted set is again an Riemann surface. We call this Riemann surface the algebraic Riemann surface determined by $P(w, z) = 0$. We call Z the set of algebraic singularities.

Suppose now we have $l \geq 2$ irreducible polynomials $P_i(w_i, z)$ with Z_i being the set of algebraic singularities of P_i. Then $Z = \bigcup_{1 \leq i \leq l} Z_i$ is a finite set. Suppose $(w_1^0, \cdots, w_l^0, z_0) \in \mathbb{C}^{l+1}$ such that $z_0 \notin Z$ and for each $1 \leq i \leq l$, $P_i(w_i^0, z_0) = 0$. Now let z run through all possible paths which start from z_0 and do not pass through points in Z, we can *simultaneously* continue all w_i to multi-valued analytic functions on $\mathbb{C} \setminus Z$. The set

$$\{(w_1(z), \cdots, w_l(z), z)\}$$

obtained in this way consist of a Riemann surface which is locally parameterized by z. The Riemann surface obtained in this way may have punctures $(w_1', \cdots, w_l', z_0')$ with $z_0' \in Z$. That is, each w_i can be extended to an analytic function in a small neighborhood of (w_i', z_0'). Let us add all such punctures to the above set. The resulted set is again a Riemann surface locally parameterized by z.

Let $f \in \mathcal{P}_\alpha^d$. The Key-Lemma 3 states one can always embed f in an algebraic Riemann surface in the parameter space

$$\underbrace{\mathbb{C}^* \times \cdots \times \mathbb{C}^*}_{(d-2) \text{ copies}}$$

such that all the periodic attracting cycles of f (if there are) are preserved with the same multipliers when moving f along the Riemann surface.

Key-Lemma 3. *Let $f \in \mathcal{P}_\alpha^d$ and $1 \leq l \leq d-3$. Suppose f has l attracting cycles with multipliers $t_1, \cdots, t_l$. Then there exist l irreducible polynomials*

$$P_i(x, y), \ 1 \leq i \leq l,$$

and a way to label the critical points of f as

$$c_1^0, \cdots, c_{d-2}^0, c_{d-1}^0 = 1,$$

such that for any $(d-1)$*-tuple*

$$X = (w_1(c_{l+1}), \cdots, w_l(c_{l+1}), c_{l+1}, c_{l+1}^0, \cdots, c_{d-2}^0, 1)$$

where w_i*,* $1 \le i \le l$*, are algebraic functions of* c_{l+1} *determined by*

$$P_i(w_i, c_{l+1}) = 0 \text{ and } w_i(c_{l+1}^0) = c_i^0,\ \ 1 \le i \le l,$$

the polynomial f_X *has* l *attracting cycles which depend holomorphically on* c_{l+1} *and has constant multiplies* $t_1, \cdots, t_l$*.*

Now consider an $f \in \Sigma_{\text{geom}}^{d,\alpha}$. Assume that f has $l+1$ periodic attracting cycles and all the critical points in the boundary of the Siegel disk satisfy some orbit relations. The Key-Lemma 4 says by sacrificing one of the attracting periodic cycles, one can embed f in an algebraic Riemann surface in the parameter space such that when the parameters vary in this algebraic Riemann surface, the other l attracting cycles and their multipliers, and all the orbit relations among the critical points on the boundary of the Siegel disk are preserved.

Key-Lemma 4. *Let* $f \in \Sigma_{\text{geom}}^{d,\alpha}$ *and* $0 \le l \le d-3$*. Suppose*

1. f *has* $l+1$ *attracting cycles with multiplies* $t_1, \cdots, t_{l+1}$ *(each of them attracts exactly one of the critical points),*
2. $f'(1) = 0$ *and* 1 *belongs to the boundary of the Siegel disk of* f*,*
3. *there are other* $d-l-3$ *critical points, say* $c_1^0, \cdots, c_{d-l-3}^0$*, on the boundary of the Siegel disk, and* $d-l-3$ *integers* $k_i \ge 1$*, such that* $f^{k_i}(1) = c_i^0$ *for* $1 \le i \le d-l-3$*.*

Then there exist a way to label the critical points of f *as*

$$c_1^0, \cdots, c_{d-2}^0, c_{d-1}^0 = 1$$

and $d-3$ *irreducible polynomials*

$$P_i(x, y),\ \ 1 \le i \le d-3,$$

such that for any $(d-1)$*-tuple*

$$X = (w_1(c_{d-2}), \cdots, w_{d-3}(c_{d-2}), c_{d-2}, 1)$$

where w_i*,* $1 \le i \le d-3$*, are algebraic functions of* c_{d-2} *determined by*

$$P_i(w_i, c_{d-2}) = 0 \text{ and } w_i(c_{d-2}^0) = c_i^0,\ \ 1 \le i \le d-3,$$

the polynomial f_X *has* l *attracting cycles which depend holomorphically on* c_{d-2} *and has constant multiplies* $t_1, \cdots, t_l$*, and moreover,* $f_X^{k_i}(1) = w_i(c_{d-2})$ *for* $1 \le i \le d-l-3$*.*

In particular, when $l = d-3$*, there is only one critical point, which is* 1*, on the boundary of the Siegel disk and thus there are no orbit relations. In this case, only* l *periodic attracting cycles are preserved and with same multipliers.*

The proofs of Key-Lemmas 3 and 4 will be given in §5. The main idea of the proofs is Euclidean algorithm. In the following we will prove the Reduced Main Theorem by assuming the four Key-Lemmas.

For $d = 2$, the Reduced Main Theorem follows from the Key-Lemma 1 directly. Suppose $d \ge 3$ and assume that the Reduced Main Theorem holds in the case that the degree is less than d. That is, there exist a pair of positive functions $\eta_0, \lambda_0 : (0, 2] \to \mathbb{R}^+$ satisfying

$$\lim_{\delta \to 0_+} \lambda_0(\delta) = \lim_{\delta \to 0_+} \eta_0(\delta) = 0,$$

such that for any pair of integers $k > m \geq 0$ and any pair of positive numbers $0 < \delta' \leq \delta$ satisfying $\delta' \leq |e^{ik\theta} - e^{im\theta}| \leq \delta$, if

$$f \in \bigcup_{\alpha \in \Theta_C^b, 2 \leq l \leq d-1} \Pi_{\text{geom}}^{\alpha, l},$$

then the inequality

$$\eta_0(\delta') \leq |f^k(1) - f^m(1)| \leq \lambda_0(\delta)$$

holds.

Take an arbitrary small $\epsilon > 0$. Let $\alpha \in \Theta_C^b$ and

$$f(z) = e^{2\pi i \alpha} z + \alpha_2 z^2 + \cdots + \alpha_d z^d$$

with $\alpha_d \neq 0$ and $f'(1) = 0$. Assume that 1 belongs to the boundary of the Siegel disk. The remaining proof is divided into three steps. In the first step, we will construct a finite chain of Riemann surfaces in the parameter space to connect f to some polynomial map g for which the boundary of the Siegel disk centered at the origin contains only the critical point 1, and moreover, there are $d-2$ periodic attracting cycles each of which attracts exactly one of the other critical points of g. The main tool used in this step is Key-Lemma 3. In the second step, we will construct a finite chain of Riemann surfaces in the parameter space to connect g to some polynomial map in the class $\Pi_{\text{geom}}^{\alpha, d}$. The main tools used in this step are Key-Lemma 2 and Key-Lemma 4. In the third step, we make the conclusion.

Step I. Assume that the number of the periodic attracting cycles of f is less than $d-2$. Otherwise, we may go to Step II directly. Let us label the critical points of f as

$$c_1^0, \cdots, c_{d-2}^0, c_{d-1}^0 = 1.$$

Fix $c_2^0, \cdots, c_{d-2}^0$ and $c_{d-1}^0 = 1$. We will repeat the following process at most $d-2$ times. Each time we will get some polynomial map which has at least one more periodic attracting cycle.

Assume that f has l attracting periodic cycles with $0 \leq l \leq d-3$. In the case $l = 0$, that is, f has no periodic attracting cycles, we just embed f into the one-parameter holomorphic family

$$h_{c_1} = f_{c_1, c_2^0, \cdots, c_{d-2}^0, 1}, \quad c_1 \in \mathbb{C}^*.$$

Otherwise, we have $1 \leq l \leq d-3$. Then by Key-Lemma 3, we can embed f into the holomorphic family

$$h_{c_{l+1}} = f_{c_1, \cdots, c_l, c_{l+1}, c_{l+2}^0, \cdots, c_{d-2}^0, 1}$$

which is locally parameterized by c_{l+1}, where all $c_i = w_i(c_{l+1}), 1 \leq i \leq l$, are algebraic functions of c_{l+1} determined respectively by the l irreducible polynomial equations

$$P_i(c_i, c_{l+1}) = 0, \ \ 1 \leq i \leq l.$$

Note that $\sigma_{k,m}(h_{c_{l+1}})$ can be regarded as a single-valued holomorphic function on a finitely many sheeted Riemann surface R branched over the c_{l+1}-punctured-plane ($\mathbb{C}^*$) with finitely many branch points, and on each sheet, there are finitely many punctures corresponding to the non-algebraic poles of the l algebraic functions $c_i = w_i(c_{l+1})$, $1 \leq i \leq l$. Recall that a point z is called an algebraic pole of an algebraic function if it is not only a pole but also a branch point. Let S_1 denote the set of all the poles (algebraic and non-algebraic) and S_2 the set of all the branch points which are not algebraic poles. Although each point in R is determined by which sheet it lies in as well as its projection to the c_{l+1}-punctured-plane, to simplify the notation, in the following discussion we just use its c_{l+1}-coordinate to denote the point when no confusion is caused. Now we have two cases.

In the first case, $h_{c_{l+1}}$ is J-stable at c_{l+1}^0. Let $U \subset R$ (In the case $l = 0$, $R = \mathbb{C}^*$) be the component in which $J_{h_{c_{l+1}}}$ moves holomorphically. Take a sequence c_{l+1}^n in U such that $\sigma_{k,m}(h_{c_{l+1}^n})$ converges to the supremum (the same argument works for the infimum for the function $\sigma_{k,m}(h_{c_{l+1}})$ does not vanish in U). By taking a subsequence we may assume that the sequence c_{l+1}^n converges to some boundary point c_{l+1}^* of U. Since the Julia set of $h_{c_{l+1}}$ moves holomorphically in U, for all the parameters in U, the boundary of the Siegel disk centered at the origin passes through the critical point 1. By the second assertion of Lemma 7.1, the critical points of $h_{c_{l+1}}$ are uniformly bounded away from the origin for all the parameters in U. Thus we have the following three subcases.

Subcase I. $c_{l+1}^* = \infty$ or $c_{l+1}^* \in S_1$. By Lemma 7.4 there is some polynomial map g of degree less than d for which the boundary of the Siegel disk contains the critical point 1 such that

$$\sigma_{k,m}(f) \leq \lim_{n\to\infty} \sigma_{k,m}(h_{c_{l+1}^n}) = \sigma_{k,m}(g) \leq \lambda_0(\delta). \tag{5}$$

Subcase II. c_{l+1}^* is a regular boundary point of U, that is, it is neither a pole nor a branch point. Then $h_{c_{l+1}}$ is not J-stable at c_{l+1}^*. Thus by McMullen-Sullivan's Theorem, one can take a $\hat{c}_{l+1}$, which can be arbitrarily close to c_{l+1}^*, such that $h_{\hat{c}_{l+1}}$ has at least one more periodic attracting cycle than $h_{c_{l+1}^*}$. Let $D_{c_{l+1}^*}$ and $D_{\hat{c}_{l+1}}$ denote respectively the Siegel disks of $h_{c_{l+1}^*}$ and $h_{\hat{c}_{l+1}}$ centered at the origin. Note that it is possible that $1 \notin \partial D_{\hat{c}_{l+1}}$. In this case, $\partial D_{\hat{c}_{l+1}}$ contains some critical point c.

Since $1 \in \partial D_{c_{l+1}^*}$ we have $\mathrm{diam}(D_{c_{l+1}^*}) \geq 1$. Since the boundary of the Siegel disk moves continuously by Lemma 7.1, by taking $\hat{c}_{l+1}$ close enough to c_{l+1}^* we can make sure that $\mathrm{diam}(D_{\hat{c}_{l+1}}) \geq 1/2$. Now from the second assertion of Lemma 7.1 we have

$$|c| \geq \mathrm{diam}(D_{\hat{c}_{l+1}})/L \geq 1/2L.$$

By taking $\hat{c}_{l+1}$ close enough to c_{l+1}^*, $\partial D_{\hat{c}_{l+1}}$, and thus c can be arbitrarily close to $\partial D_{c_{l+1}^*}$. Thus for the given $\epsilon > 0$, by taking $\hat{c}_{l+1}$ close enough to c_{l+1}^*, we can make sure that there exist a pair of integers $k' > m' > 0$ such that $k' - m' = k - m$ and

$$\sigma_{k,m}(h_{c_{l+1}^*}) = |h_{c_{l+1}^*}^k(1) - h_{c_{l+1}^*}^m(1)| < |h_{\hat{c}_{l+1}}^{k'}(c) - h_{\hat{c}_{l+1}}^{m'}(c)| + \epsilon.$$

Let

$$h(z) = c^{-1} h_{\hat{c}_{l+1}}(cz).$$

Then the boundary of the Siegel disk of h centered at the origin passes through the critical point 1 and

$$|h_{\hat{c}_{l+1}}^{k'}(c) - h_{\hat{c}_{l+1}}^{m'}(c)| < 2L \cdot \sigma_{k',m'}(h).$$

We finally get

$$\sigma_{k,m}(f) \le \lim_{n\to\infty} \sigma_{k,m}(h_{c_{l+1}^n}) = \sigma_{k,m}(h_{c_{l+1}^*}) < 2L\cdot \sigma_{k',m'}(h) + \epsilon. \tag{6}$$

Subcase III. $c_{l+1}^* \in S_2$. We may assume that c_{l+1}^* is not an isolated boundary point. This is because otherwise, in a small punctured neighborhood of c_{l+1}^*, by using a coordinate transformation $c_{l+1} = \zeta^j + c_{l+1}^*$ where $j \ge 2$ is the branch degree at c_{l+1}^*,

$$F(\zeta) = \sigma_{k,m}(h_{c_{l+1}})(\zeta^j + c_{l+1}^*)$$

is a holomorphic function of ζ in an open punctured neighborhood of the origin. As $\zeta \to 0$, $F(\zeta)$ converges to $\sigma_{k,m}(h_{c_{l+1}^*})$ which is a finite complex number. Thus $\zeta = 0$ is a removable singularity of $F(\zeta)$. Since $F(\zeta)$ obtains maximal at $\zeta = 0$, it follows that $F(\zeta)$ is a constant in a small neighborhood of the origin. It follows that $\sigma_{k,m}(h_{c_{l+1}})$ is a constant in U. Since there are only finitely many branched points and poles, we can assume that $c_{l+1}^* = \infty$ or is a regular boundary point of U. Hence we go back to Subcase I or Subcase II and hence either (5) or (6) holds.

Now assume that c_{l+1}^* is not an isolated point in ∂U. Since there are only finitely many poles and branch points, we can take a regular boundary point of U, say $\tilde{c}_{l+1}$, in an arbitrarily small neighborhood of c_{l+1}^*. Since the Julia set of $h_{c_{l+1}}$ moves holomorphically in U, the critical point 1 belongs to the boundaries of the Siegel disks for all the parameters in U. In particular, for both $h_{\tilde{c}_{l+1}}$ and $h_{c_{l+1}^*}$, the boundaries of the Siegel disks centered at the origin contain the critical point 1. For the given $\epsilon > 0$, by Lemma 7.4, we can take $\tilde{c}_{l+1}$ close to c_{l+1}^* enough so that

$$\sigma_{k,m}(h_{c_{l+1}^*}) < \sigma_{k,m}(h_{\tilde{c}_{l+1}}) + \epsilon.$$

Since $\tilde{c}_{l+1}$ is a regular boundary point of U, as in Subcase II, we can find $h \in \mathcal{P}_\alpha^d$ which has at least one more periodic attracting cycle than $h_{\tilde{c}_{l+1}}$ and for which the boundary of the Siegel disk centered at the origin contains the critical point 1, and moreover, there are two integers $k' > m' \ge 0$ with $k' - m' = k - m$ such that

$$\sigma_{k,m}(h_{\tilde{c}_{l+1}}) < 2L\cdot \sigma_{k',m'}(h) + \epsilon.$$

So finally we have

$$\sigma_{k,m}(f) \le \sigma_{k,m}(h_{c_{l+1}^*}) < \sigma_{k,m}(h_{\tilde{c}_{l+1}}) + \epsilon < 2L\cdot \sigma_{k',m'}(h) + 2\epsilon. \tag{7}$$

In the second case, $h_{c_{l+1}}$ is J-unstable at c_{l+1}^0. Then by regarding c_{l+1}^0 as c_{l+1}^* and using the same argument as in the Subcase II of the first case, we can find $h \in \mathcal{P}_\alpha^d$ which has at least one more periodic attracting cycle than $h_{c_{l+1}^0}$ and for which the boundary of the Siegel disk centered at the origin contains the critical point 1, and moreover, there are two integers $k' > m' \ge 0$ with $k' - m' = k - m$ such that (6) holds.

Now we repeat the above process on h. Since each time the number of the attracting periodic cycles is increased by at least one, there are two possibilities. The first one is we get an inequality like (5) at some step, and therefore get

$$\sigma_{k,m}(f) \le 2L\cdot(2L\cdot(\cdots(2L\cdot\lambda_0(\delta) + 2\epsilon) + \cdots) + 2\epsilon) + 2\epsilon, \tag{8}$$

where the number of the recursive steps is not greater than $d-3$. The second one is we get a pair of integers $k' > m' \ge 0$ with $k' - m' = k - m$ and a polynomial map $g \in \mathcal{P}_\alpha^d$ which has $d-2$ periodic attracting cycles, each of which attracts a critical point, and moreover, the boundary of the Siegel disk centered at the origin contains the critical point 1, such that

$$\sigma_{k,m}(f) \leq 2L \cdot (2L \cdot (\cdots (2L \cdot \sigma_{k',m'}(g) + 2\epsilon) + \cdots) + 2\epsilon) + 2\epsilon, \tag{9}$$

where the number of the recursive steps is not greater than $d-2$.

If we get (8) we go to **Step III** directly. So assume that we get the above polynomial map g satisfying (9).

Step II. Note that g has $d-2$ attracting cycles and each of them attracts exactly one critical point, and the boundary of the Siegel disk centered at the origin contains the only one critical point 1. We will repeat the following process $d-2$ times.

Suppose $0 \leq l \leq d-3$ and $g \in \mathcal{P}_\alpha^d$ has $l+1$ attracting cycles, and there are $d-l-2$ critical points on the boundary of the Siegel disk, say $c_1^0, \cdots, c_{d-l-3}^0$, and 1. By Key-Lemma 2, we can get a polynomial $\tilde{g} \in \mathcal{P}_\alpha^d$ arbitrarily close to g such that all the critical points on the boundary of the Siegel disk satisfy orbit relations, that is, $\tilde{g}^{k_i}(1) = c_i$ for $1 \leq i \leq d-l-3$; and moreover, $\tilde{g}$ has $l+1$ attracting cycles with the same multipliers as g. Because $\tilde{g}$ can be arbitrarily close to g, we may assume that

$$\sigma_{k',m'}(g) < \sigma_{k',m'}(\tilde{g}) + \epsilon. \tag{10}$$

Note that we start from the case $l = d-3$. In this case the point 1 is the only critical point on the boundary of the Siegel disk, and thus there is no orbit relations among critical points on the boundary of the Siegel disk.

By Key-Lemma 4, we can embed $\tilde{g}$ in a holomorphic family

$$h_{c_{d-2}} = f_{c_1, \cdots, c_{d-3}, c_{d-2}, 1}$$

such that $h_{c_{d-2}}$ has l attracting cycles with constant multipliers $t_1, \cdots, t_l$, and all the orbit relations are preserved, that is, $h_{c_{d-2}}^{k_i}(1) = c_i$, $1 \leq i \leq d-l-3$, where c_i, $1 \leq i \leq d-3$, are determined by $d-3$ irreducible polynomial equations $P_i(c_i, c_{d-2}) = 0$.

Again, $\sigma_{k',m'}(h_{c_{d-2}})$ is a holomorphic function defined on a finitely many sheeted Riemann surface R branched over the c_{d-2}-plane. As in Step I let S_1 denote the set of all the (algebraic) poles and S_2 the set of all the branch points which are not (algebraic) poles. Let $S = S_1 \cup S_2$. Then S is a finite set.

Let $\Sigma \subset R \setminus S$ be the subset which contains all those points for which the boundary of the Siegel disk of $h_{c_{d-2}}$ contains exactly those $d-l-2$ critical points, $c_1, \cdots, c_{d-l-3}$, 1, which satisfy the orbit relations. We claim that Σ is an open set. Let us prove the claim. Suppose $c_{d-2} \in \Sigma$. Let $D_{c_{d-2}}$ denote the Siegel disk of $h_{c_{d-2}}$ centered at the origin. Then all the critical points of $h_{c_{d-2}}$, which do not belong to $\partial D_{c_{d-2}}$, have a positive distance from $\partial D_{c_{d-2}}$. By Lemma 7.3, in a small open neighborhood of c_{d-2}, all these critical points are still bounded away from $\partial D_{c_{d-2}}$. It suffices to show that the critical points, which satisfy the orbit relations, still belong to $\partial D_{c_{d-2}}$. Let us prove this by contradiction. Note that except 1, there are exactly $d-l-3$ critical points, $c_1, \cdots, c_{d-3}$ on the boundary of the Siegel disk, and $d-l-3$ integers, $1 \leq k_1 < \cdots < k_{d-l-3}$ such that $f^{k_i}(1) = c_i$, and moreover, there are l attracting cycles with multipliers $t_1, \cdots, t_l$. If the claim were not true, then there would be a sequence $c_{d-2}^n \to c_{d-2}$ such that for $h_{c_{d-2}^n}$, at least one of the $d-l-2$ critical points, 1, $c_1^n, \cdots, c_{d-l-3}^n$, does not belong to $\partial D_{c_{d-2}^n}$. For the convenience, let us denote $c_0^n = 1$. By taking a subsequence, we may assume that there is a an $0 \leq i \leq d-l-3$ such that $c_i^n \notin \partial D_{c_{d-2}^n}$, and $c_j^n \in \partial D_{c_{d-2}^n}$, for all $i+1 \le j \le d-l-3$. Consider the map

$$F_{c_{d-2}} = h_{c_{d-2}}^{k_{i+1}-k_i}.$$

The map $F_{c_{d-2}}$ maps c_i to c_{i+1} and the local degree of $F_{c_{d-2}}$ at c_i is equal to that of $h_{c_{d-2}}$ at c_i, which is two. This means there is a pair of Jordan domains A and B containing c_i and c_{i+1} respectively such that $F_{c_{d-2}} : A \to B$ is a branched covering map of degree two. Then for all c_{d-2}^n close to c_{d-2} enough, there is a pair of Jordan domains A_n and B_n with $A_n \to A$ and $B_n \to B$ such that $F_{c_{d-2}^n} : A_n \to B_n$ is a branched covering map of degree two. But on the other hand, when c_{d-2}^n is close to c_{d-2}, c_{i+1}^n is close to c_{i+1} and thus belongs to B_n and c_i^n is close to c_i and thus belongs to A_n. Let $z_n \in \partial D_{c_{d-2}^n}$ be the point such that $F_{c_{d-2}^n}(z_n) = c_{i+1}^n$. As $c_{d-2}^n \to c_{d-2}$, we must have $z_n \to c_i$. Thus as n is large enough, z_n belongs to A_n also. This implies that the preimage of c_{i+1}^n in A_n under the map $F_{c_{d-2}^n}$, , counting by multiplicities, is at least three. This is a contradiction. Thus Σ is an open set and the claim has been proved.

Let U denote the component of Σ which contains the point c_{d-2}^0. Since for all $c_{d-2} \in U$, 1 belongs to the boundary of the Siegel disk on which $h_{c_{d-2}}$ is qc-conjugate to the irrational rotation R_α, $\sigma_{k',m'}(h_{c_{d-2}})$ does not vanish in U. So both the maximal and minimal principles can apply to $\sigma_{k',m'}(h_{c_{d-2}})$ in U. Take a sequence c_{d-2}^n in U such that $\sigma_{k',m'}(h_{c_{d-2}^n})$ converges to its supremum (The same argument works for the infimum). By taking a subsequence, we may assume that c_{d-2}^n converges to a point $c_{d-2}^* \in \partial U$.

Since for all $c_{d-2} \in U$, the boundary of the Siegel disk of $h_{c_{d-2}}$ centered at the origin passes through the critical point 1, by the second assertion of Lemma 7.1, all critical points of $h_{c_{d-2}}$ are uniformly bounded away from the origin. Thus we have the following subcases.

Subcase I. $c_{d-2}^* = \infty$ or $c_{d-2}^* \in S_1$. By Lemma 7.4 there is some polynomial map h of degree less than d for which the boundary of the Siegel disk contains the critical point 1 such that

$$\sigma_{k',m'}(\tilde{g}) \le \lim_{n\to\infty} \sigma_{k',m'}(h_{c_{d-2}^n}) = \sigma_{k',m'}(h) \le \lambda_0(\delta). \tag{11}$$

Subcase II. c_{d-2}^* is a regular boundary point of U. We thus have

$$\sigma_{k',m'}(\tilde{g}) \le \sigma_{k',m'}(h_{c_{d-2}^*}). \tag{12}$$

We claim that $h_{c_{d-2}^*}$ has exactly one more critical point on the boundary of the Siegel disk, that is, there are $d-l-1$ critical points on $\partial D_{c_{d-2}^*}$. This is because the boundary of the Siegel disk moves continuously by Lemma 7.3, since all the $d-l-2$ critical points $c_1^n, \cdots, c_{d-l-3}^n$, 1 belong to $\partial D_{c_{d-2}^n}$, all the $d-l-2$ critical points $c_1^*, \cdots, c_{d-l-3}^*$, 1 must belong to $h_{c_{d-2}^*}$ also. If there is no more critical point on $\partial D_{c_{d-2}^*}$, then $c_{d-2}^* \in \Sigma$. Because Σ is open, there exists an open disk neighborhood of c_{d-2}^*, say V in Σ. Since $c_{d-2}^n \to c_{d-2}^*$ and $c_{d-2}^n \in U$, it follows that $U \cup V$ is a connected and open subset of Σ. This contradicts the assumption that U is a connected component of Σ. So there must be at least one more critical point in $\partial D_{c_{d-2}^*}$. Since $h_{c_{d-2}^*}$ has l attracting cycles which attract at least l critical points, it follows that $\partial D_{c_{d-2}^*}$ contains exactly one more critical point. The claim has been proved.

Subcase III. $c_{d-2}^* \in S_2$. Then as in the Subcase III in Step I, by changing the coordinate $c_{d-2} = c_{d-2}^* + \zeta^j$ for some appropriate integer j we may assume that c_{d-2}^* is not an isolated boundary point of U, for otherwise, we will go back to Subcases I or II. Thus we can choose a regular boundary point $\hat{c}_{d-2}$ arbitrarily close to c^*_{d-2}. By Lemma 7.4 we an take $\hat{c}_{d-2}$ close to c^*_{d-2} enough such that

$$\sigma_{k',m'}(\tilde{g}) \leq \sigma_{k',m'}(h_{c^*_{d-2}}) < \sigma_{k',m'}(h_{\hat{c}_{d-2}}) + \epsilon. \tag{13}$$

As in the Subcase II, $\partial D_{\hat{c}_{d-2}}$ contains exactly one more critical point, that is, $d-l-1$ critical points, and $h_{\hat{c}_{d-2}}$ has l attracting periodic cycles, each of which attracts exactly one critical point.

Now we can repeat the above process. Since each time the number of the critical points on the boundary of the Siegel disk is increased by one, after at most $d-2$ steps, we can either get an inequality like (11) at some step or we finally get a Siegel polynomial map $h \in \Pi^{\alpha,d}_{\text{geom}}$. From (10)-(13) we have either

$$\sigma_{k',m'}(g) \leq 2L \cdot (2L \cdot (\cdots (2L \cdot \lambda_0(\delta) + 2\epsilon) + \cdots) + 2\epsilon) + 2\epsilon, \tag{14}$$

where the number of the recursive steps is not more than $d-2$ or

$$\sigma_{k',m'}(g) \leq 2L \cdot (2L \cdot (\cdots (2L \cdot \sigma_{k',m'}(h) + 2\epsilon) + \cdots) + 2\epsilon) + 2\epsilon, \tag{15}$$

where the number of the recursive steps is not more than $d-2$ and $h \in \Pi^{\alpha,d}_{\text{geom}}$. From Key-Lemma 1, we have $\sigma_{k',m'}(h) \leq \lambda_1(\delta)$. Thus (15) can be rewritten into

$$\sigma_{k',m'}(g) \leq 2L \cdot (2L \cdot (\cdots (2L \cdot \lambda_1(\delta) + 2\epsilon) + \cdots) + 2\epsilon) + 2\epsilon, \tag{16}$$

where the number of recursive steps is not more than $d-2$.

From (8) and(14), (9) and (16) we have in all the cases

$$\sigma_{k,m}(f) \leq 2L \cdot (2L \cdot (\cdots (2L \cdot \max\{\lambda_0(\delta) + \lambda_1(\delta)\} + 2\epsilon) + \cdots) + 2\epsilon) + 2\epsilon, \tag{17}$$

where the number of the recursive steps is $2d$ times. Since $\epsilon > 0$ is arbitrary, by letting $\epsilon \to 0$, we have

$$\sigma_{k,m}(f) \leq (2L)^{2d} \cdot \max\{\lambda_0(\delta), \lambda_1(\delta)\}.$$

In the same way we can prove

$$\sigma_{k,m}(f) \geq (2L)^{-2d} \cdot \min\{\eta_0(\delta), \eta_1(\delta)\}.$$

Now define $\lambda, \eta : (0,2] \to (0, +\infty)$ by setting

$$\lambda(x) = (2L)^{2d} \cdot \max\{\lambda_0(x), \lambda_1(x)\} \quad \text{and} \quad \eta(x) = (2L)^{-2d} \cdot \min\{\eta_0(x), \eta_1(x)\}.$$

The proof of the Reduced Main Theorem is complete.

3. Proof of Proposition 2.1

Let $F : \mathbb{C} \to \mathbb{C}$ be a continuous map. For any pair of integers $k > m \geq 0$, define

$$\sigma_{k,m}(F) = |F^k(1) - F^m(1)|. \tag{18}$$

Suppose $\lambda, \eta : (0,1] \to \mathbb{R}^+$ are a pair of positive functions such that

$$\lim_{\delta \to 0_+} \lambda(\delta) = \lim_{\delta \to 0_+} \eta(\delta) = 0. \tag{19}$$

Lemma 3.1. *Suppose for any pair of integers $k > m \geq 0$ and any pair of positive numbers $0 < \delta' < \delta$ satisfying*

$$\delta' < |e^{2k\pi i\theta} - e^{2m\pi i\theta}| < \delta,$$

we have

$$\eta(\delta') \leq |F^k(1) - F^m(1)| \leq \lambda(\delta).$$

Then

$$\Gamma = \overline{\{F^k(1)\}_{k=0}^{\infty}}$$

is a Jordan curve. Moreover, $F : \Gamma \to \Gamma$ is a topological circle homeomorphism with rotation number θ.

Proof. Let $\mathbb{T}$ denote the unit circle. Then $X = \{e^{2k\pi i\theta}\}_{k\geq 1}$ is a dense subset of $\mathbb{T}$. Define a map $\phi : X \to \mathbb{C}$ by setting $\phi(e^{2k\pi i\theta}) = F^k(1)$ for every $k \geq 0$. Then $\phi : X \to \mathbb{C}$ is uniformly continuous by assumption. Thus ϕ can be uniquely extended to a continuous map from $\mathbb{T}$ to $\mathbb{C}$. Let us still denote the map by ϕ. We claim that ϕ is injective.

Let us prove it by contradiction. Assume that $\phi(x) = \phi(y)$ for some $x \neq y \in \mathbb{T}$. Let $\delta' = \frac{1}{2}|x - y|$. Since X is dense in $\mathbb{T}$ and $\phi : \mathbb{T} \to \mathbb{C}$ is continuous, we have two integers $k > m \geq 0$ such that

(1) $|e^{2k\pi i\theta} - x| < |x - y|/4$,
(2) $|e^{2m\pi i\theta} - y| < |x - y|/4$,
(3) $|\phi(e^{2k\pi i\theta}) - \phi(x)| < \eta(\delta')/2$,
(4) $|\phi(e^{2m\pi i\theta}) - \phi(y)| < \eta(\delta')/2$.

From (1) and (2) we have $|e^{2k\pi i\theta} - e^{2m\pi i\theta}| > |x - y|/2 = \delta'$. From the assumption, we have $|\phi(e^{2k\pi i\theta}) - \phi(e^{2m\pi i\theta})| > \eta(\delta')$. But from (3), (4) and $\phi(x) = \phi(y)$, we have $|\phi(e^{2k\pi i\theta}) - \phi(e^{2m\pi i\theta})| < \eta(\delta')$. This is a contradiction. This implies that $\phi : \mathbb{T} \to \mathbb{C}$ is injective. Thus

$$\overline{\{F^k(1)\}_{k=0}^{\infty}} = \overline{\phi(X)} = \phi(\mathbb{T})$$

is a Jordan curve. This proves the first assertion.

To prove the second assertion, note that

$$F \circ \phi = \phi \circ R_\theta$$

holds on X. Since X is dense on $\mathbb{T}$, ϕ is continuous on $\mathbb{T}$ and F is continuous on $\mathbb{C}$, the above equation still holds on $\mathbb{T}$. Since ϕ is injective on $\mathbb{T}$, it follows that

$$\phi^{-1} \circ F \circ \phi = R_\theta$$

holds on $\mathbb{T}$. This proves the second assertion.

□

This implies Proposition 2.1.

4. PROOF OF KEY-LEMMA 1

4.1. **Uniform real bounds.** We shall identify the unit circle with $\mathbb{T} = \mathbb{R}/\mathbb{Z}$ and give $\mathbb{T}$ the induced orientation. Let $f : \mathbb{T} \to \mathbb{T}$ be a homeomorphism. Let (a, b, c, d) be a quadruple with $a < b < c < d < a + 1$. Define the cross-ratio

$$[a, b, c, d] = \frac{b - a}{c - a}\frac{d - c}{d - b}$$

and the cross-ratio distortion

$$D(a,b,c,d,f) = \frac{[f(a),f(b),f(c),f(d)]}{[a,b,c,d]}.$$

Let $d \geq 2$ be an integer. Let

$$\mathcal{H}_d = \big\{ g(z) = \lambda z^d \prod_{i=1}^{d-1} \frac{1-\overline{a}_i z}{z-a_i}, 0<|a_i|<1, g|_{\mathbb{T}} \text{ is a homeomorphism} \big\}.$$

Lemma 4.1 (Herman, [9])**.** *There is a $0 < C(d) < \infty$ depending only on d such that for any quadruple (a,b,c,d), any $g \in \mathcal{H}_d$ and any integer $m \geq 1$, if all the intervals in the collection*

$$\{f^i(I)\}_{i=0}^{m-1}, \text{ where } I=(a,b),$$

are disjoint with each other, then

$$D(a,b,c,d,f^m) < C(d).$$

Lemma 4.2 (Uniform power law)**.** *There exist constants $\nu, C' > 1$ such that for any $g \in \mathcal{H}_d$, if $c \in \mathbb{T}$ is a critical point of g, then for any $x, y \in \mathbb{T}$ with $|x-c| \leq |y-c|$, we have*

$$\left|\frac{g(x)-g(c)}{g(y)-g(c)}\right| \leq C' \cdot \left|\frac{x-c}{y-c}\right|^{\nu}.$$

Proof. Let $I = [c,y]$ and $J = [c,x]$. By Lemma 15 of [**?**], when restricted to an open neighborhood of $\mathbb{T}$, the maps in $\mathcal{H}_d$ consist of a normal family. Thus there exist a pair of positive numbers $0 < m < M$ which depend only on d and an open neighborhoods U of $\mathbb{T}$ such that

$$g'(z) = \phi(z) \cdot (z-c)^l \cdot \prod_{i\in\Lambda}(z-c_i)^{l_i} \cdot \prod_{i\in\Theta}(z-c_i)^{l_i}$$

where

1. $m \leq |\phi(z)| \leq M$ for all $z \in U$,
2. $\{c\} \cup \{c_i, i \in \Lambda \cup \Theta\} = \{z \in U \mid g'(z) = 0\}$,
3. $\text{dist}(c_i, c) \geq 2|I|$ for all $c_i \in \Lambda$ and $\text{dist}(c_i, c) < 2|I|$ for all $c_i \in \Theta$.

Note that except 0 and ∞ g has $2(d-1)$ critical points, counting by multiplicities. So there exist constants $0 < \kappa(d), \eta(d) < 1$ depending only on d and a sub-interval $I_1 \subset I$ such that

- i $|I_1| > \eta(d) \cdot |I|$,
- ii $\text{dist}(c, I_1) > \eta(d) \cdot |I|$,
- iii for any $c_i \in \Theta$, $\text{dist}(c_i, I_1) > \kappa(d) \cdot |I|$.

For any $c_i \in \Theta$ and $z \in J$, it is clear that $\text{dist}(c_i, z) \leq \text{dist}(c_i, c) + \text{dist}(c, z)$. Since $\text{dist}(c_i, c) < 2|I|$ and $\text{dist}(c, z) \leq |J| < |I|$, we have

$$\sup_{z\in J} \text{dist}(c_i, z) < 3|I|. \tag{20}$$

For any $c_i \in \Lambda$ and $z \in J$, it is clear that $\text{dist}(c_i, z) < \text{dist}(c_i, I_1) + \text{dist}(z, I_1)$. Since $\text{dist}(z, I_1) \leq \text{dist}(z, c) + \text{dist}(c, I_1) \leq |J| + |I| < 2|I|$ and $\text{dist}(c_i, I_1) \geq \text{dist}(c_i, c) - \text{dist}(c, I_1) > \text{dist}(c_i, c) - |I| \geq |I|$, we have

$$\sup_{z\in J} \text{dist}(c_i, z) < \text{dist}(c_i, I_1) + \sup_{z\in J} \text{dist}(z, I_1) < \text{dist}(c_i, I_1) + 2|I| < 3\text{dist}(c_i, I_1). \tag{21}$$

Now from the intermediate value theorem we have $\xi \in I_1$ and $\zeta \in J$ such that

$$|g(I)| \geq |g(I_1)| = |g'(\xi)| \cdot |I_1| = |\phi(\xi)| \cdot |\xi - c|^l \cdot \prod_{i \in \Lambda} |\xi - c_i|^{l_i} \cdot \prod_{i \in \Theta} |\xi - c_i|^{l_i} \cdot |I_1|,$$

and

$$|g(J)| = |g'(\zeta)| \cdot |J| = |\phi(\zeta)| \cdot |\zeta - c|^l \cdot \prod_{i \in \Lambda} |\zeta - c_i|^{l_i} \cdot \prod_{i \in \Theta} |\zeta - c_i|^{l_i} \cdot |J|.$$

Since g is of degree of $2d-1$ we have $2 \leq l, l_i \leq 2d-2$ for all $i \in \Theta \cup \Lambda$. This, together with (i-iii) and (20-21), implies

$$\frac{|g(J)|}{|g(I)|} \leq \frac{M}{m} \cdot \frac{|J|^l}{\eta(d)^l \cdot |I|^l} \cdot 3^{2d-2} \cdot \frac{3^{2d-2}}{\kappa(d)^{2d-2}} \cdot \frac{1}{\eta(d)}.$$

Since $2 \leq l \leq 2d-2$ the lemma follows by taking $\nu = 2$ and

$$C' = \frac{M \cdot 3^{4d-4}}{m \cdot \eta(d)^{2d-1} \cdot \kappa(d)^{2d-2}}.$$

□

Let $B \in \mathcal{H}_d$ such that the rotation number of $B|\mathbb{T} : \mathbb{T} \to \mathbb{T}$ is an irrational number and the point 1 is a critical point of B. It is necessary that the local degree of B at 1 is odd and not less than 3. We use backward notation. Let x_i denote the point in $\mathbb{T}$ such that $B^i(x_i) = 1$. Let

$$I_n = [1, x_{q_n}] \text{ and } I_{n+1} = [1, x_{q_{n+1}}].$$

For $i \geq 0$, let I_n^i denote the subintervals of $\mathbb{T}$ such that $B^i(I_n^i) = I_n$. Then the collections of the intervals

$$\Pi_n(B) = \{I_n^i,\ 0 \leq i \leq q_{n+1} - 1;\ \ I_{n+1}^i,\ 0 \leq i \leq q_n - 1\} \tag{22}$$

form a partition of $\mathbb{T}$. We call $\Pi_n(B)$ the dynamical partition of level n. For $K > 1$ and $I, J \subset \mathbb{T}$, we say I and J are K-commensurable if $|J|/K < |I| < K|J|$.

Theorem 4.1 (Herman-Swiatek's uniform real bounds)**.** *There is a $1 < C(d) < \infty$ depending only on d such that for any $B \in \mathcal{H}_d$ for which $B|\mathbb{T} : \mathbb{T} \to \mathbb{T}$ is a circle homeomorphism with rotation number θ and has a critical point at 1, we have*

1. *the two intervals $[x, B^{q_n}(x)]$ and $[x, B^{-q_n}(x)]$ are $C(d)$-commensurable for any $x \in \mathbb{T}$ and $n \geq 1$.*
2. *any two adjacent intervals in the dynamical partition of level n are $C(d)$-commensurable for any $n \geq 1$.*

Proof. For the proof, see Proposition 3.3 and Corollary 3.6 of [15]. We only need to notice that the constants obtained in [15] only depend on those in the two assumptions of Hypothesis 1 of [15] and in our situation, these two assumptions are satisfied because of Lemmas 4.1 and 4.2. □

4.2. **The family $\mathcal{S}_d$.** Let $d \geq 2$ be an integer and $\alpha \in \Theta_C$ be an irrational number of bounded type. Let $f : \mathbb{T} \to \mathbb{T}$ be a circle homeomorphism and $\phi : \mathbb{T} \to \mathbb{T}$ be a homeomophism such that

$$f = \phi^{-1} \circ R_\alpha \circ \phi.$$

We define the *dynamical angle* between two points x and y in $\mathbb{T}$ to be the angle between $\phi(x)$ and $\phi(y)$ anticlockwise. Recall that $\Pi_{\text{geom}}^{\alpha,d}$ denote the class of all polynomial maps of degree d which has a critical point at 1 and a Siegel disk centered at the origin and of rotation number α such that the boundary of the Siegel disk contains all the finite critical points. The next lemma provides the premodels for all the Siegel polynomial maps in the class $\Pi_{\text{geom}}^{\alpha,d}$.

Lemma 4.3. *For any tuple $(\alpha_1, \cdots, \alpha_{d-1})$ with $0 \leq \alpha_i < 2\pi$ and $\sum_{i=1}^{d-1} \alpha_i = 2\pi$, there exists a $B \in \mathcal{H}_d$ such that*

1. *there are exactly $d-1$ critical points $c_1 = 1, c_2, \cdots, c_{d-1}$ in $\mathbb{T}$, ordered anticlockwise and counted by multiplicities,*
2. *the angle from c_i to c_{i+1} anticlockwise is α_i for $1 \leq i \leq d-1$ (we identify c_1 with c_d),*
3. *the rotation number of $B|\mathbb{T} : \mathbb{T} \to \mathbb{T}$ is α.*

Proof. It is is a direct consequence of Theorem 2.1. □

Let $\mathcal{S}_d \subset \mathcal{H}_d$ denote the class of all the Blaschke products guaranteed by Lemma 4.3. In §6, we shall see that each $f \in \Pi_{\text{geom}}^{\alpha,d}$ is uniquely determined by the group of angles $(\alpha_1, \cdots, \alpha_{d-1})$ formed by the $d-1$ critical points on the boundary of the Siegel disk, and thus for any $f \in \Pi_{\text{geom}}^{\alpha,d}$, there exists a $B \in \mathcal{S}_d$ such that f is obtained from B through a qc surgery.

4.3. **Uniform saddle node geometry.** Take $B \in \mathcal{S}_d$. Then there is a circle homeomorphism $h : \mathbb{T} \to \mathbb{T}$ such that

$$B|\mathbb{T} = h^{-1} \circ R_\alpha \circ h.$$

Here $0 < \alpha < 1$ is a fixed irrational number in Θ_C. The aim of this subsection is to show that the non-linearity of h is characterized by uniform saddle node geometry which depends only on d and C. This is the most subtle part in the whole paper: the uniform estimate relies essentially on the compactness property of the family $\mathcal{S}_d$, and does not hold for the corresponding family for rational maps.

First let us give some notations. For $i \geq 0$, let $x_i \in \mathbb{T}$ be the point such that $B^i(x_i) = 1$. Let p_n/q_n be the n-th convergent of α. For each $n \geq 0$, define

$$\mathcal{Q}_n = \{x_i \mid 0 \leq i < q_n\}.$$

Then $\mathcal{Q}_0 = \{1\}$. The following proposition is summarized from §6 of [17] where the circle homeomorphism is induced by the Douady-Ghys Blaschke model which contains exactly one (double) critical point at 1. Since it only involves the combinatorial information about the rotation number and is independent of the number of the critical points in $\mathbb{T}$, it still holds for $B \in \mathcal{S}_d$. Recall that $\Pi_n(B)$ is the collection of intervals in the dynamical partition of level n defined in (22).

Proposition 4.1 (cf. §6 of [17])**.** Let $0 \le j < k < q_n$. Then x_j and x_k are adjacent in $\mathcal{Q}_n$ if and only if $k = j + q_{n-1}$ and $0 \le j < q_n - q_{n-1}$, or $k = j + q_n - q_{n-1}$ and $0 \le j < q_{n-1}$. In the former case we have

$$[x_k, x_j] \cap \mathcal{Q}_{n+1} = \{x_k, x_{k+q_n}, x_{k+2q_n}, \cdots, x_{k+(a_{n+1}-1)q_n}, x_j\},$$

and in the later case we have

$$[x_j, x_k] \cap \mathcal{Q}_{n+1} = \{x_j, x_{j+q_n}, x_{j+2q_n}, \cdots, x_{j+a_{n+1}q_n}, x_k\}.$$

Moreover, each interval in $\mathbb{T} \setminus \mathcal{Q}_n$ is either one interval in $\Pi_n(B)$, or the union of two adjacent intervals in $\Pi_{n-1}(B)$. In particular, any two adjacent intervals in $\mathbb{T} \setminus \mathcal{Q}_n$ are K-commensurable with $K > 1$ being some constant depending only on d.

The last assertion of the above proposition is implies by the second assertion of Lemma 4.1 that any two adjacent intervals in $\Pi_{n-1}(B)$ are $C(d)$-commensurable with $C(d) > 1$ being some constant depending only on d.

When $d = 2$, B is exactly the Douady-Ghys Blaschke model considered in [17]. In this case the point 1 is the unique critical point of B in $\mathbb{T}$. It is known that for every interval component I of $\mathbb{T} \setminus \mathcal{Q}_n$, the map

$$B^{q_n} : I \to f^{q_n}(I)$$

is a diffeomorphism (cf. §6 of [17]).

For $d > 2$, there are $(d-1)$ critical points in $\mathbb{T}$, counting by multiplicities. If all these critical points collide into one critical point at 1, then for any interval component I of $\mathbb{T} \setminus \mathcal{Q}_n$, the map $B^{q_n} : I \to I$ is still a diffeomorphism. Otherwise, there are $1 \le d' \le d-2$ distinct critical points other than 1. Let us denote them by $c_i, 1 \le i \le d'$ and denote 1 by c_0. For any integer $k \ge 0$, let $c_i^k \in \mathbb{T}$ denote the point such that $B^k(c_i^k) = c_i$. Let

$$\mathcal{Q}_n^i = \{c_i^k \mid \ 0 \le k < q_n\},\ 1 \le i \le d',\ n \ge 1.$$

Then $\mathcal{Q}_n^0 = \mathcal{Q}_n$.

Lemma 4.4. *Let $n \ge 1$. Then for each $1 \le i \le d'$, we have*

1. *in the case that $k = j + q_{n-1}$ and $0 \le j < q_n - q_{n-1}$, the interval (x_k, x_j) contains at most one point in $\mathcal{Q}_n^i$,*
2. *in the case that $k = j + q_n - q_{n-1}$ and $0 \le j < q_{n-1}$, the interval (x_j, x_k) contains at most two points in $\mathcal{Q}_n^i$.*

Proof. Fix an $1 \le i \le d'$. By replacing $\mathcal{Q}_n$ by $\mathcal{Q}_n^i$ in Proposition 4.1, it follows that each component of $\mathbb{T} \setminus \mathcal{Q}_n^i$ either has the form $[c_i^m, c_i^l]$, where $m = l + q_{n-1}$ and $0 \le l \le q_n - q_{n-1}$, or has the form $[c_i^l, c_i^m]$ where $m = l + q_n - q_{n-1}$ and $0 \le l < q_{n-1}$.

For $k = j + q_{n-1}$ and $0 < j < q_n - q_{n-1}$, It is clear that (x_k, x_j) can not contain any interval of the form $[c_i^m, c_i^l]$ with $m = l + q_{n-1}$ and $0 \le l < q_n - q_{n-1}$. By the property of closest returns, the interval (x_k, x_j) can not contain any interval of the form $[c_i^l, c_i^m]$ with $m = l + q_n - q_{n-1}$ and $0 \le l < q_{n-1}$. In this case, we must have $q_n - q_{n-1} > q_{n-1}$. This implies that (x_k, x_j) contains at most one point in $\mathcal{Q}_n^i$. This proves the first assertion.

For $k = j + q_n - q_{n-1}$ and $0 \le j < q_{n-1}$, it is clear that the interval (x_j, x_k) can contain at most one interval of the form $[c_i^m, c_i^l]$ with $m = l + q_{n-1}$ and $0 \le l \le q_n - q_{n-1}$ and can not contain any interval with the form $[c_i^l, c_i^m]$ with $m = l + q_n - q_{n-1}$ and $0 \le l < q_{n-1}$. This implies that (x_j, x_k) can contain at most two points in $\mathcal{Q}_n^i$. This proves the second assertion. □

Let I be an interval component of $\mathbb{T}\backslash \mathcal{Q}_n$. Then the points in $\mathcal{Q}_{n+1}$ divide I into finitely many sub-intervals. By Theorem 4.1, any two such sub-intervals are K-commensurable for some constant $K > 1$ depending only on d. When the number of such sub-intervals are large, however, the ones which lie in the middle position are very small compared to the ones near the end. This is the so called "saddle-node" geometry. More precisely,

Lemma 4.5 (cf. Theorem 6.6 of [17]). *Let $d = 2$. Then there is a universal constant $K > 1$ such that for any interval component I of $\mathbb{T} \setminus \mathcal{Q}_n$, if the points in $\mathcal{Q}_{n+1}$ divide I into sub-intervals*

$$I_1, \cdots, I_m,$$

then we have

$$\frac{1}{K} \cdot \frac{|I|}{\min\{|k|, |k-m+1|\}^2} < |I_k| < K \cdot \frac{|I|}{\min\{|k|, |k-m+1|\}^2}$$

The aim of this subsection is to general the above lemma to the case that $d > 2$.

Lemma 4.6 (Uniform Saddle Node Geometry). *There is a $K > 1$ depending only on d such that for any $B \in \mathcal{S}_d$, if I is a component of $\mathbb{T} \setminus \mathcal{Q}_n$, then for any component J of*

$$I \setminus \bigcup_{i=1}^{d-2} \mathcal{Q}_n^i$$

such that J contains at least one interval component of $I \setminus \mathcal{Q}_{n+1}$, we have

$$\frac{1}{K} \cdot |I| < |J| \leq |I|.$$

Moreover, if

$$J_1, \cdots, J_m$$

denote all the interval components of $I \setminus \mathcal{Q}_{n+1}$ contained in J, then we have

$$\frac{1}{K} \cdot \frac{|J|}{\min\{|k|, |k-m+1|\}^2} < |J_k| < K \cdot \frac{|J|}{\min\{|k|, |k-m+1|\}^2}.$$

The proof of Lemma 4.6 uses Yoccoz's almost parabolic lemma, which is the same tool as in the proof of Lemma 4.5. Before we state this lemma let us introduce some terminologies first. Let n a be a positive integer and let $I_1, \cdots, I_n$ be consecutive intervals on the line or circle. By an almost parabolic map of length n and fundamental domains $I_1, \cdots, I_n$, we mean a negative-Schwarzian diffeomorphism

$$f : I_1 \cup \cdots I_n \to I_2 \cup \cdots \cup I_{n+1}$$

such that $f(I_j) = I_{j+1}$. The basic geometric estimate on almost parabolic maps is due to J.-C. Yoccoz.

Lemma 4.7 (Yoccoz's Lemma). *Let $f : I \to f(I)$ be an almost parabolic map of length n and fundamental domains $I_j, 1 \leq j \leq n$. If $|I_1| \geq \sigma \cdot |I|$ and $|I_n| \geq \sigma \cdot |I|$ for some $\sigma > 0$, then*

$$\frac{1}{C_\sigma} \frac{|I|}{\min\{j, n-j\}^2} \leq |I_j| \leq C_\sigma \frac{|I|}{\min\{j, n-j\}^2}$$

where $C_\sigma > 1$ is a constant depending only on σ.

The first assertion of Lemma 4.6 is implied by the following lemma.

Lemma 4.8. *Let I and J be the intervals in Lemma 4.6. Let $\mathcal{Q}_n^0 = \mathcal{Q}_n$. Then for any interval component S of $I \setminus \mathcal{Q}_{n+1}$, if*

$$\overline{S} \cap \bigcup_{i=0}^{d-2} \mathcal{Q}_n^i \neq \emptyset,$$

then $|S| > |I|/C$ where $C > 1$ is some constant depending only on d.

Proof. The argument is quite standard. By Proposition 4.1 we have two cases: either $I = (x_k, x_j)$ when $k = j + q_{n-1}$ and $0 \le j < q_n - q_{n-1}$, or $I = (x_j, x_k)$ when $k = j + q_n - q_{n-1}$ and $0 \le j < q_{n-1}$. By Proposition 4.1 again, each of these two cases has two subcases. In the first case, S either has the form $(x_{k+lq_n}, x_{k+(l+1)q_n})$ for some $0 \le l \le a_{n+1} - 2$, or has the form $(x_{k+(a_{n+1}-1)q_n}, x_j)$, and in the second case, S either has the form $(x_{j+lq_n}, x_{j+(l+1)q_n})$ for some $0 \le l \le a_{n+1} - 1$ or has the form $(x_{j+a_{n+1}q_n}, x_k)$. Since the proofs of all these four subcases are similar to each other, let us only deal with the first subcase. With a minor changes of the proof, the reader shall easily prove the remaining three subcases.

Now we can suppose $I = (x_k, x_j)$ where $k = j + q_{n-1}$ and $0 \le j < q_n - q_{n-1}$ and $S = (x_{k+lq_n}, x_{k+(l+1)q_n})$ for some $0 \le l \le a_{n+1} - 2$. Since $\overline{S} \cap \bigcup_{i=0}^{d-2} \mathcal{Q}_n^i \neq \emptyset$, there is a least integer $0 \le t \le q_n - 1$ and a point $x \in \overline{S}$ such that $B^t(x) = c$ where $c \in \mathbb{T}$ is some critical point of B.

Let $y_0 = c$ and for $i > 0$, let y_i denote the point in $\mathbb{T}$ such that $B^i(y_i) = c$. Consider the following two group of intervals

$$\text{I. } [y_0, y_{q_n}], [y_{q_n}, y_{q_n - q_{n-1}}], [y_{q_n - q_{n-1}}, y_{q_n - 2q_{n-1}}]$$

and

$$\text{II. } [y_{2q_{n-1}}, y_{q_{n-1}}], [y_{q_{n-1}}, y_0], [y_0, y_{q_n}].$$

These intervals belong to the collection of the intervals of the dynamical partition of $\mathbb{T}$ of level $n-1$ with respect to the critical point c. By Theorem 4.1, for each of the two group, the three intervals are $C(d)$-commensurable with $C(d) > 1$ is some constant depending only on d. Thus the cross ratios of both $(y_0, y_{q_n}, y_{q_n - q_{n-1}}, y_{q_n - 2q_{n-1}})$ and $(y_{2q_{n-1}}, y_{q_{n-1}}, y_0, y_{q_n})$ have a lower bound $\kappa(d) > 0$ depending only on d. Pull back the two group of intervals by B^{-t}. We get the following two group of intervals

$$\text{III. } [x, B^{-q_n}(x)], [B^{-q_n}(x), B^{-q_n + q_{n-1}}(x)], [B^{-q_n + q_{n-1}}(x), B^{-q_n + 2q_{n-1}}(x)]$$

and

$$\text{IV. } [B^{2q_{n-1}}(x), B^{q_{n-1}}(x)], [B^{q_{n-1}}(x), x], [x, B^{q_n}(x)].$$

Since $0 \le t < q_n$, the pull backs of each interval in I and II by $B^i, i = 0, 1, \cdots, t$, are disjoint. Thus the intersection multiplicity of the pull backs of each of the two groups I and II is not greater than $3(d-1)$.

Now by Lemma 4.1 and the compactness property of the family $\mathcal{S}_d$, it follows that the cross ratios of both

$$(x, B^{-q_n}(x), B^{-q_n + q_{n-1}}(x), B^{-q_n + 2q_{n-1}}(x))$$

and

$$(B^{2q_{n-1}}(x), B^{q_{n-1}}(x), x, B^{q_n}(x))$$

have a positive lower bound $\eta(d) > 0$ with $\eta(d) > 0$ being some constant depending only on d. This implies that

$$|[x, B^{-q_n}(x)]| > \lambda(d)\cdot|[B^{-q_n}(x), B^{-q_n+q_{n-1}}(x)]| \text{ and } |[x, B^{-q_n}(x)]| > \lambda(d)\cdot|[B^{q_{n-1}}(x), x]| \tag{23}$$

with $\lambda(d) > 0$ being some constant depending only on d.

Since $I = (x_k, x_j) = (B^{-q_{n-1}}(x_j), x_j)$ and $x \in I$, we have

$$I \subset [B^{q_{n-1}}(x), x] \cup [x, B^{-q_n}(x)] \cup [B^{-q_n}(x), B^{-q_n+q_{n-1}}(x)].$$

This, together with (23), implies that

$$|[x, B^{-q_n}(x)]| > \frac{\lambda(d)}{2} \cdot |I|. \tag{24}$$

By the first assertion we have

$$[x_{k+(l+1)q_n}, x_{k+(l+2)q_n}]| < C(d) \cdot |[x_{k+lq_n}, x_{k+(l+1)q_n}]|$$

where $C(d) > 1$ is some constant depending only on d. Since

$$x \in S = [x_{k+lq_n}, x_{k+(l+1)q_n}],$$

we have

$$[x, B^{-q_n}(x)] \subset \overline{S} \cup [x_{k+lq_n}, x_{k+(l+1)q_n}]. \tag{25}$$

From (24) and (25) we have

$$|S| > \frac{1}{1 + C(d)} \cdot |[x, B^{-q_n}(x)]| > \frac{1}{1 + C(d)} \cdot \frac{\lambda(d)}{2} \cdot |I|.$$

□

Let $J_1, \cdots, J_m$ be the intervals in Lemma 4.6. From Proposition 4.1 it is known J_1, J_2, and J_3 are K-commensurable for some constant $K > 1$ depending only on d. Let us consider the diffeomorphism

$$B^{q_n} : J_3 \cup \cdots \cup J_m \to J_2 \cup \cdots \cup J_{m-1}.$$

From Lemma 4.7, we see that to prove the second assertion of Lemma 4.6, it suffices to check two conditions. The first one is to show that the two boundary sub-intervals, that is, J_3 and J_m, are uniformly commensurable with the whole interval J. The second one is to show that B_n^q has negative Schwarze derivative on $J_3 \cup \cdots J_m$. The following lemma implies the first condition.

Lemma 4.9. *Let J and $J_1, \cdots, J_m$ be intervals. Then there exist $\sigma > 0$ depending only on d such that $|J_1| > \sigma \cdot |J|$ and $|J_m| > \sigma \cdot |J|$.*

Proof. Let us prove the first inequality only. The second one can be proved by the same argument. There are two cases. In the first case J_1 has a common boundary point with J. Then the boundary point must be a point in $\bigcup_{i=1}^{d-2} \mathcal{Q}_n^i$. By Lemma 4.8, we have $|J_1| > |I|/C > |J|/C$ for some $C > 1$ depending only on d. In the second case, J_1 is adjacent to an interval component of $I \setminus \mathcal{Q}_{n+1}$, say S, which contains a boundary point of J. By Proposition 4.1, S and J_1 are K-commensurable for some $K > 1$ depending only on d. On the other hand, by Lemma 4.8 we have $|S| > |I|/C > |J|/C$ for some $C > 1$ depending only on d. This implies that $|J_1| > (KC)^{-1} \cdot |J|$. This proves the lemma. □

It remains to prove that B^{q_n} has negative Schwarz derivative on $J_3 \cup \cdots \cup J_m$. Here when we talk about the Schwarz derivatives of B and its iterations, we regard $\mathbb{T}$ as $\mathbb{R}/\mathbb{Z}$ and $B : \mathbb{T} \to \mathbb{T}$ as its lift $f : \mathbb{R} \to \mathbb{R}$. In this way, B is real in $\mathbb{R}$ and analytic in a strip neighborhood of $\mathbb{R}$, and moreover, $B(x+1) = B(x)+1$. To simplify the notation we still use B to denote this map if no confusion arises.

Lemma 4.10. *There is a $C > 1$ depending only on d such that for any x and y in $J_3 \cup \cdots \cup J_m$ and all $1 \le k \le q_n$, we have*

$$C^{-1} < \frac{DB^k(x)}{DB^k(y)} < C.$$

Proof. It is known that the map $B^{-q_n} : J_2 \cup \cdots \cup J_{m-1} \to J_3 \cup \cdots \cup J_m$ is a diffeomorphism. Let U be the Euclidean disk such that ∂U is orthogonal to $\mathbb{T}$ and passes through the two end points of the arc

$$J_1 \cup \cdots \cup J_m.$$

Then U contains $J_2 \cup \cdots \cup J_{m-1}$ in its interior, and moreover, since J_1 and J_m are K-commensurable with $J_1 \cup \cdots \cup J_m$ and $K > 1$ is a constant depending only on d, it follows that the modulus of the annulus $U \setminus \overline{J_2 \cup \cdots \cup J_{m-1}}$ has a positive lower bound depending only on d. Since U does not intersect the post-critical set of B, B^{-q_n} can be holomorphically extended to a univalent function on U. Let $V = B^{-q_n}(U)$. Then the modulus of $V \setminus \overline{J_3 \cup \cdots \cup J_m}$ is equal to that of $U \setminus \overline{J_2 \cup \cdots \cup J_{m-1}}$ and thus has a positive lower bound depending only on d. It is clear that for every $1 \le k \le q_n$, the map B^k is univalent in V. The lemma then follows from the Koebe's distortion theorem. ☐

Lemma 4.11. *There is an $N \ge 1$ such that for all $n \ge N$ and all $B \in \mathcal{S}_d$, if J_i, $1 \le i \le m$, are the intervals in Lemma 4.6, then $S(B^{q_n})(z) < 0$ for all $z \in J_3 \cup \cdots \cup J_m$.*

Proof. Recall that we regard B as a holomorphic function defined in a strip neighborhood of $\mathbb{R}$ such that $B : \mathbb{R} \to \mathbb{R}$ is a homeomorphism and $B(x+1) = B(x) + 1$. Since $\mathcal{S}_d$ is compact, one can take a $\xi > 0$ such that all $B \in \mathcal{S}_d$ are holomorphic in

$$S = \{z = x + iy \mid -\xi < y < \xi\}.$$

In particular, $B'(x)$ is a periodic function with period 1. Since B' has at most $d-1$ distinct zeros in each interval $[x, x+1)$, there is a $0 < \kappa < 1$ such that for each $B \in \mathcal{S}_d$, we can find a $t \in [0,1)$ such that all zeros of B' in $[t-\kappa, t+1+\kappa]$ belong to $(t, t+1)$. Let $c_1, \cdots, c_{d-1}$ denote all the zeros of B' in $(t, t+1)$. Let $U = \{x+iy \mid t-\kappa < x < t+1+\kappa, -\xi < y < \xi\}$. For $z \in U$, let

$$B'(z) = g(z) \cdot \Pi_{1 \le i \le d-1}(z - c_i)^2.$$

Then g is a holomorphic function defined in U. Let $V = \{x + iy \mid t - \kappa/2 < x < t+1+\kappa/2, -\xi/2 < y < \xi/2\}$. Since $\mathcal{S}_d$ is compact, by a compactness argument it follows that there is a $0 < \eta < 1$ depending only on d such that for all $z \in V$, we have

i. $|g(z)| \ge \eta$,
ii. $|g'(z)| < 1/\eta$,
iii. $|g''(z)| < 1/\eta$.

Let $x \in (t - \kappa/2, t + 1 + \kappa/2)$ and $x \ne c_i$, $1 \le i \le d-1$. Let

$$P(x) = \Pi_{1 \le i \le d-1}(x - c_i).$$

Then $B'(x) = P(x)^2 \cdot g(x)$. By direct calculations we have

$$\begin{aligned} B''(x) &= 2P(x)P'(x)g(x) + P(x)^2 g'(x) \\ &= 2P^2(x)\Sigma(x)g(x) + P^2(x)g'(x) = 2P^2(x)\Sigma(x)g(x) + P^2(x)g'(x) \\ &= P^2(x)(2\Sigma(x)g(x) + g'(x)) \end{aligned}$$

where

$$\Sigma(x) = P'(x)/P(x) = \sum_{1 \le i \le d-1} \frac{1}{x - c_i},$$

and

$$B'''(x) = 2P^2(x)\Sigma(x)(2\Sigma(x)g(x) + g'(x)) + P^2(x)(-2\sigma(x)g(x) + 2\Sigma(x)g'(x) + g''(x))$$

where

$$\sigma(x) = -\Sigma'(x) = \sum_{1 \le i \le d-1} \frac{1}{|x - c_i|^2}.$$

Then

$$\frac{B'''(x)}{B'(x)} = 4\Sigma^2(x) + 4\Sigma(x)\frac{g'(x)}{g(x)} - 2\sigma(x) + \frac{g''(x)}{g(x)}$$

and

$$\frac{B''(x)}{B'(x)} = 2\Sigma(x) + \frac{g'(x)}{g(x)}.$$

From

$$SB(x) = \frac{B'''(x)}{B'(x)} - \frac{3}{2}(\frac{B''(x)}{B'(x)})^2,$$

we finally have

$$SB(x) = -2\Sigma^2(x) - 2\Sigma(x)\frac{g'(x)}{g(x)} - 2\sigma(x) + \frac{g''(x)}{g(x)} - \frac{3}{2}(\frac{g''(x)}{g(x)})^2. \tag{26}$$

Let $\delta = \min_{1 \le i \le d-1} |x - c_i|$. From the properties (i), (ii) and (iii), it follows that there is a $0 < C < \infty$ such that

1. $-2\Sigma^2(x) < 0$,
2. $|-2\Sigma(x)\frac{g'(x)}{g(x)}| \le \frac{C}{\delta}$,
3. $-2\sigma(x) \le -\frac{2}{\delta^2}$,
4. $|\frac{g''(x)}{g(x)} - \frac{3}{2}(\frac{g''(x)}{g(x)})^2| < C$.

All of these implies that

$$SB(x) < -\frac{1}{\delta^2}$$

provided that $\delta = \min_{1 \le i \le d-1}$ is small enough. Since $B'(x)$ is periodic with period one, this is true for all $x \in \mathbb{R}$ provided that $\delta = \mathrm{dist}(x, \Omega_B)$ is small enough where Ω_B is the set of all critical points in $\mathbb{R}$.

Now let us prove the lemma. Let $\tilde{J}$ denote one of the lifts of $J_3 \cup \cdots \cup J_m$ to $\mathbb{R}$. Let $\tilde{J}_i = B^i(\tilde{J})$ for $i \ge 1$. Let $x \in \tilde{J}$ be an arbitrary point. Let us consider the sum

$$SB^{q_n}(x) \cdot |\tilde{J}|^2 = \sum_{j=0}^{q_n - 1} SB(B^j(x))(DB^j(x))^2 \cdot |\tilde{J}|^2. \tag{27}$$

By Lemma 4.10, it follows that

$$K^{-1} \cdot |\tilde{J}_j|^2 < (DB^j(x))^2 \cdot |\tilde{J}|^2 < K \cdot |\tilde{J}_j|^2$$

where $K > 1$ is some constant depending only on d. Take an $\epsilon > 0$ small such that the above inequality holds.

Let $U = \{x \mid \operatorname{dist}(x, \Omega_B) < \epsilon\}$ and $V = \{x \mid \operatorname{dist}(x, \Omega_B) > \epsilon/2\}$. Take an N such that $\tilde{J}_j$ is either contained in U or contained in V. We split the sum in (27) into Σ_1 and Σ_2: Σ_1 consists of all the terms such that $\tilde{J}_i$ is contained in U, and Σ_2 contains all the other terms.

Note that all the terms in Σ_1 is negative. Moreover, by Theorem 4.1, there is a $\tilde{J}_j$ such that $\operatorname{dist}(\tilde{J}_j, \Omega_B)$ is K-commensurable with $|\tilde{J}_j|$ with $K > 1$ depending only on d. For such j we have

$$SB(B^j(x)) < -\frac{1}{K^2 \cdot |\tilde{J}_j|^2}.$$

This implies that there is a $\eta > 0$ such that

$$\Sigma_1 < -\eta$$

provided that $n \geq N$.

On the other hand, for all $x \in V$, $|SB(x)| < C$ for some $0 < C < \infty$ depending only on d. Since the projections of all $\tilde{J}_j$, $0 \leq j \leq q_n - 1$ to $\mathbb{T}$ are disjoint, we have

$$\sum_{j=0}^{q_n-1} |\tilde{J}_j| < 1.$$

By taking N large enough, we can make sure that $|\tilde{J}_j| < (KC)^{-1} \cdot \eta$. Then for all $n \geq N$, we have

$$\Sigma_2 < C \cdot \sum_{j=0}^{q_n-1} K \cdot |\tilde{J}_j|^2 \leq \eta.$$

This completes the proof of the lemma. □

Now apply Lemma 4.7 to the diffeomorphism $B^{q_n} : J_3 \cup \cdots \cup J_m \to J_2 \cup \cdots \cup J_{m-1}$. The two conditions in Lemma 4.7 are satisfied by Lemmas 4.9 and 4.11. Lemma 4.6 now follows from Lemma 4.7.

4.4. **Proof of Lemma 2.1.** The aim of this subsection is to prove Lemma 2.1. We will prove that there exist uniform constants $M, \alpha > 0$ and $0 < \epsilon_0 < 1$ such that for any $B \in \mathcal{S}_d$, the circle homeomorphism $h_B : \mathbb{T} \to \mathbb{T}$ induced by B has a David extension $H : \Delta \to \Delta$ which is a (M, α, ϵ_0)-David homeomorphism. The basic idea used here is the same as that used in [17]: For each of the two circle homeomorphisms $h : \mathbb{T} \to \mathbb{T}$ and $R_\alpha : \mathbb{T} \to \mathbb{T}$, we give a way to divide Δ into countably many polygons. For each pair of corresponding polygons, we construct a qc homeomorphism between them so that the restriction of the homeomorphism to each edge of the polygon is linear. The map H is obtained by gluing all these polygon homeomorphisms along the edges. Compared with the construction in [17], a slight difference arises here. In Petersen-Zakeri's proof, the bottom side of each polygon consists of one single piece of saddle node geometry, while in our case, the bottom side of each polygon consists of several pieces of saddle node geometry. Our idea is to divide each polygon into finitely many pieces so that each new one has one single saddle node geometry. In this way we can use the idea in [17]. The following lemma, which is essentially the Lemma 6.5 in [17], is a fundamental block in this construction.

Lemma 4.12 (cf. Lemma 6.5, [17]). *Let P and Q be two unit squares. Let*

$$X = \{x_1, x_2, \cdots, x_m\} \text{ and } Y = \{y_1, y_2, \cdots, y_m\}$$

be two partitions of the two bottom sides of P and Q, respectively. Then P and Q become into two polygons by regrading the previous vertices of P and Q, and the points in X and Y as the new vertices of P and Q respectively. Suppose the partition X has C_0-bounded saddle-node geometry for some $C_0 > 1$, that is,

$$\frac{1}{C_0}\frac{|x_1 - x_m|}{\min\{i, m-i\}^2} \le |x_i - x_{i+1}| \le C_0 \frac{|x_1 - x_m|}{\min\{i, m-i\}^2}, \ 1 \le i \le m-1,$$

and Y has the C_1-bounded linear geometry for some $C_1 > 1$, that is,

$$\frac{1}{C_1 \cdot m} \cdot |y_1 - y_m| \le |y_i - y_{i+1}| \le \frac{C_1}{m} \cdot |y_1 - y_m|, \ 1 \le i \le m-1.$$

Then there is a K-qc homeomorphism $F : P \to Q$ such that when restricted to the corresponding edges, F is a linear map and

$$K < C \cdot (1 + (\log m)^2)$$

where $C > 1$ is a constant depending only on C_0 and C_1.

Recall a continuous curve segment is called a polyline if it connects finitely many distinct points $x_1, \cdots, x_m$ successively through straight segments.

Definition 4.1. *Suppose $C > 1$ and L is a polyline which connects $x_1, \cdots, x_m$. We say L has a C-bounded uniform saddle node geometry if*

$$\frac{1}{C}\frac{|x_1 - x_m|}{\min\{i, m-i\}^2} \le |x_i - x_{i+1}| \le C \frac{|x_1 - x_m|}{\min\{i, m-i\}^2}, \ 1 \le i \le m-1$$

and C-bounded linear geometry if

$$\frac{1}{C \cdot m} \cdot |x_1 - x_m| \le |x_i - x_{i+1}| \le \frac{C}{m} \cdot |x_1 - x_m|, \ 1 \le i \le m-1.$$

To construct a David extension $H : \Delta \to \Delta$ of the circle mapping $h : \mathbb{T} \to \mathbb{T}$ with uniform parameters $M, \alpha > 0$ and $0 < \epsilon_0 < 1$, we need a generalized version of Lemma 4.12.

Lemma 4.13. *Let P and Q be two unit squares. Let*

$$X = \{x_1, x_2, \cdots, x_m\} \text{ and } Y = \{y_1, y_2, \cdots, y_m\}$$

be two partitions of the two bottom sides of P and Q, respectively. Then P and Q become into two polygons by regrading the previous vertices of P and Q, the points in X and Y as the new vertices of P and Q respectively. Let $l \ge 1$.

Suppose the partition X consists of l pieces all of which have C_0-bounded saddle-node geometry for some $C_0 > 1$, that is,

1. *there exist*
$$1 = m_0 < m_1 < m_2 < \cdots < m_{l-1} < m_l = m$$
such that
$$|x_0 - x_m|/C_0 \le |x_{m_{i-1}} - x_{m_i}| \le |x_0 - x_m|, \ 1 \le i \le l,$$
and
2. *for* $0 \le i \le m-1$ *and* $m_i \le j < m_{i+1}$,
$$\frac{1}{C_0}\frac{|x_{m_i} - x_{m_{i+1}}|}{\min\{j - m_i + 1, m_{i+1} - j\}^2} \le |x_j - x_{j+1}| \le C_0 \frac{|x_{m_i} - x_{m_{i+1}}|}{\min\{j - m_i + 1, m_{i+1} - j\}^2}.$$

Suppose in addition that Y *has* C_1*-bounded linear geometry for some* $C_1 > 1$*, that is,*
$$\frac{1}{C_1 \cdot m} \cdot |y_1 - y_m| \le |y_j - y_{j+1}| \le \frac{C_1}{m} \cdot |y_1 - y_m|, \ 1 \le j \le m-1.$$
Then there is a K*-qc homeomorphism* $F : P \to Q$ *such that when restricted to the corresponding edges,* F *is a linear map and*
$$K < C \cdot (1 + (\log m)^2)$$
where $C > 1$ *is a constant depending only on* C_0*,* C_1 *and* l*.*

Proof. Let A and B denote the two vertices of the bottom side of P. Let A' and B' denote the two corresponding vertices of the bottom side of Q.

If $l = 1$, then the lemma is implied by Lemma 4.12.

Suppose $l \ge 2$ and the lemma holds for $l - 1$. Let us prove the lemma holds for l. Then either $m_1 - m_0 \ge m_l - m_{l-1}$ or $m_1 - m_0 \le m_l - m_{l-1}$. Let us assume that $m_1 - m_0 \ge m_l - m_{l-1}$. The case that $m_1 - m_0 \le m_l - m_{l-1}$ can be treated in the same way. Let
$$n = \left[\frac{m_l - m_0}{m_l - m_{l-1}}\right] + 4$$
where $[\,\cdot\,]$ denote the integer part of a number. Then $n \ge 6$.

Claim: There exist $K_1, C_2 > 1$ depending only on C_0, and $C_3, K_2 > 1$ depending only on C_1, and two group of points $x_1', \cdots, x_n'$ and $y_1', \cdots, y_n'$, such that

1. $x_1' = x_1 = A$, $x_{n-3}' = x_{m_1}$, $x_n' = x_{m_2} = B$,
2. x_j' lies in the interior of P for all $2 \le j \le n-4$ and $j = n-2, n-1$,
3. $y_1' = y_1 = A'$, $y_{n-3}' = y_{m_1}$, $y_n' = y_{m_2} = B'$,
4. y_j' lies in the interior of Q for all $2 \le j \le n-4$ and $j = n-2, n-1$,

and moreover, if L and L' are two polylines connecting $x_1', \cdots, x_n'$, and $y_1', \cdots, y_n'$ respectively, then

5. L has C_2-bounded saddle-node geometry,
6. Let L_1 be the part of L connecting A and x_{m_1} and L_2 be the remaining part of L. There exist a polyline S between L_1 and the straight segment $[A, x_{m_1}]$ which consists of three straight segments and connects A and x_{m_1} such that the following properties hold. P is divided by L and S into four polygons P_1, P_2, P_3, P_4, where P_1 is the top one, P_2 is the one at the right-lower corner, P_3 is the one bounded by S and L_1, P_4 is the one bounded by S and $[A, x_{m_1}]$. Moreover, for each $i = 1, 2, 3,$ 4, there is a K_1-qc homeomorphism ϕ_i mapping

P_i to a polygon which comes from a standard square with the bottom side being divided into finitely many pieces satisfying either C_0 or C_2-bounded saddle-node geometry(cf. Lemma 4.12 for the description of such polygon). More precisely, for P_1, L is mapped to the bottom side which satisfies C_2-bounded saddle node geometry; for P_2, $[x_{m_1}, B]$ is mapped to the bottom side which satisfies C_0-bounded saddle-node geometry; for P_3, L_1 is mapped to the bottom side which satisfies C_2-bounded saddle-node geometry; for P_4, $[A, x_{m_1}]$ is mapped to the bottom side which consists of $(l-1)$ pieces each of which satisfies C_0-bounded saddle-node geometry. For each P_i, $1 \leq i \leq 4$, the map is linear on each edge of P_i and maps each edge of P_i to the corresponding edge of the polygon.

7. L' has C_3-bounded linear geometry,
8. Let L_1' be the part of L' connecting A' and y_{m_1} and L_2' be the remaining part of L'. There exists a polyline segment S' between L_1' and the straight segment $[A', y_{m_1}]$ which consists of three straight segments and connects A' and y_{m_1} such that the following properties hold. Q is divided by S' and L' into four polygons Q_1, Q_2, Q_3, Q_4, where Q_1 is the top one, Q_2 is the one at the right-lower corner, Q_3 is the one bounded by S' and L_1', Q_4 is the one bounded by S' and $[A', y_{m_1}]$. Moreover, for each i, there is a K_2-qc homeomorphism ψ_i mapping Q_i to a polygon which comes from a standard square with the bottom side being divided into finitely many pieces satisfying either C_1 or C_3-bounded linear geometry (cf. Lemma 4.12 for a description of such polygons). More precisely, for Q_1, L' is mapped to the bottom side satisfying C_3-bounded linear geometry, for Q_2, $[y_{m_1} B']$ is mapped to the bottom side satisfying C_1-bounded linear geometry, for Q_3, L_1' is mapped to the bottom side satisfying C_3-bounded linear geometry, for Q_4, $[A', y_{m_1}]$ is mapped to the bottom side which consists of $(l-1)$ pieces each of which satisfies C_1-bounded linear geometry. For each Q_i, $1 \leq i \leq 4$, the map is linear on each edge of Q_i and maps each edge of Q_i to the corresponding edge of the polygon.

Note that $n \leq m$. By Lemma 4.12 and the induction assumption, there exists a $C' > 1$ depending only on C_0, C_1, C_2, C_3, l and thus only on C_0, C_1 and l (since C_2 and C_3 depend respectively on C_0 and C_1) such that for each $i = 1, 2, 3, 4$, there is a K-qc homeomorphism σ_i mapping $\phi_i(P_i)$ to $\psi_i(Q_i)$ and satisfying the following properties.

1. $K < C' \cdot (1 + (\log m)^2)$,
2. each edge of $\phi_i(P_i)$ is linearly mapped to the corresponding edge of $\psi_i(Q_i)$.

Note that P_i is mapped to Q_i by $\psi_i^{-1} \circ \sigma_i \circ \phi_i$ such that each edge of P_i is linearly mapped to the corresponding edge of Q_i. Since ϕ_i is K_1-qc and ψ_i is K_2-qc, thus $\psi_i^{-1} \circ \sigma_i \circ \phi_i$ is $K_1 \cdot K_2 \cdot C'$-qc. Since K_1 and K_2 depends respectively on C_0 and C_1 and C' depends on C_0, C_1 and l, it follows that $C = K_1 \cdot K_2 \cdot C'$ depends only on C_0, C_1 and l. We can now glue the maps $\psi_i^{-1} \circ \sigma_i \circ \phi_i$ along the edges of P_i and get a qc map $F : P \to Q$ with qc constant bounded by $C \cdot (1 + (\log m)^2)$. This proves the lemma by assuming the claim.

Now let us prove the Claim. Let us first describe how to choose the points x_i', $i = 0, \cdots, n$. Let $a = |[A, x_{m_{l-1}}]|$ and $b = |[x_{m_{l-1}}, B]|$. Let $x_0' = x_0 = A$, $x_{n-3}' = x_{m_{l-1}}$ and $x_n' = B$. Let x_1' be the point in the interior of P such that the length of $[x_0', x_1']$ is equal to $a/2$, and the angle formed by $[x_0', x_1']$ and $[A, B]$ is $\pi/3$. Let x_{n-4}' be the point in the interior of P such that the length of $[x_{n-4}', x_{n-3}']|$ is equal to $a/2$ and the angle formed



FIGURE 1. Divide P and Q into four polygons with one side with saddle-node and linear geometry respectively

by $[x'_{n-4}, x'_{n-3}]|$ and $[x_{m_{l-1}}, A]$ is equal to $\pi/3$. Then the straight segment $[x'_1, x'_{n-4}]$ has length $a/2$. There is an obvious way to insert points $x'_2, \cdots, x'_{n-5}$ in the interior of $[x'_1, x'_{n-4}]$ so that the horizontal polyline

$$[x'_1, \cdots, x'_{n-4}]$$

satisfies Θ-bounded saddle-node geometry with $\Theta > 1$ being some universal constant.

Now let x'_{n-2} be the point in the interior of P such that the length of $[x'_{n-3}, x'_{n-2}]$ is equal to $b/2$ and the angle formed by $[x'_{n-3}, x'_{n-2}]$ and $[x'_{n-3}, B]$ is $\pi/3$. Let x'_{n-1} be the point in the interior of P such that the length of $[x'_n, x'_{n-1}]$ is equal to $b/2$ and the angle formed by $[x'_n, x'_{n-1}]$ and $[B, A]$ is equal to $\pi/3$. Then the length of the horizontal straight segment $[x'_{n-2}, x'_{n-1}]$ is $b/2$. Since a and b are C_0-commensurable, and $[x'_1, \cdots, x'_{n-4}]$ satisfies Θ-bounded saddle-node geometry, it follows that the polyline

$$L = [x'_0, \cdots, x'_n]$$

satisfies C_2-bounded saddle node geometry with $C_2 > 1$ depends only on C_0.

Now let us describe how to construct the polyline S. Let $s_0 = A$ and $s_3 = x_{m_{l-1}}$. Let s_1 be the point in the interior of P such that the length of $[s_0, s_1]$ is equal to $a/3$ and the angle formed by $[s_0, s_1]$ and $[A, B]$ is $\pi/4$. Let s_2 be the point in the interior of P such that the length of $[s_2, s_3]$ is equal to $a/3$ and the angle formed by $[s_2, s_3]$ and $[x_{m_{l-1}}, A]$ is $\pi/3$. Let S be the polyline which connects s_0, s_1, s_2 and s_3.

Let L_1 bet he part of L connecting x'_0 and $x_{m_{l-1}}$ and L_2 be the remaining part of L. Then from the construction we see that L and S divide P into four polygons P_i, $1 \le i \le 4$: P_1 is the top one; P_2 is the one bounded by L_2 and $[x_{m_{l-1}}, B]$; P_3 is the one bounded by L_1 and S; and P_4 is the one bounded by S and $[A, x_{m_{l-1}}]$. Each of these polygons has four sides, the three of which are straight segments and the last one is a polyline with either C_0 or C_2-bounded saddle node geometry. Here $C_2 > 1$ is some constant depending only on C_0. Since the geometry of each P_i is well controlled, it is not difficult to construct the desired qc homeomorphisms ϕ_i which maps each P_i to a standard square with the bottom side being divided into finitely many pieces satisfying either C_0 or C_2-bounded saddle node geometry.

Let us now describe how to construct the polylines L' and Q'. Let $a' = |[A', y_{m_{l-1}}]|$ and $b' = |[y_{m_{l-1}}, B']|$. By assumption it follows that

$$a' \geq \frac{(n-5)b'}{C_1^2}.$$

Let $y'_0 = y_0 = A'$, $y'_{n-3} = y_{m_{l-1}}$ and $y'_n = B'$. Let $n_1 = [n/3]$ and $n_2 = [2n/3]$. Let y'_{n_1} be the point in the interior of Q such that the length of $[y'_0, y'_{n_1}]$ is equal to $a'/2$, and the angle formed by $[y'_0, y'_{n_1}]$ and $[A', B']$ is $\pi/3$. Let y'_{n_2} be the point in the interior of Q such that the length of $[y'_{n_2}, y'_{n-3}]|$ is equal to $a'/2$ and the angle formed by $[y'_{n_2}, y'_{n-3}]|$ and $[y_{m_{l-1}}, A']$ is equal to $\pi/3$. Then the straight segment $[y'_1, y'_{n_2}]$ has length $a'/2$. Now we insert points $y'_1, \cdots, y'_{n_1-1}$ in the interior of $[y'_0, y'_{n_1}]$, and $y'_{n_1+1}, \cdots, y'_{n_2-1}$ in the interior of $[y'_{n_1}, y'_{n_2}]$, and $y'_{n_2+1}, \cdots, y'_{n-4}$ in the interior of $[y_{n_2}, y_{n-3}]$, so that

$$|[y'_i, y'_{i+1}]| = |[y'_0, y'_{n_1}]/(n_1), \ \ 0 \leq i \leq n_1 - 1,$$

$$|[y'_i, y'_{i+1}]| = |[y'_{n_1}, y'_{n_2}]/(n_2 - n_1), \ \ n_1 \leq i \leq n_2 - 1,$$

and

$$|[y'_i, y'_{i+1}]| = |[y'_{n_2}, y'_{n-3}]/(n - 3 - n_2), \ \ n_2 \leq i \leq n - 4.$$

Now let y'_{n-2} be the point in the interior of Q such that the length of $[y'_{n-3}, y'_{n-2}]$ is equal to $b'/2$ and the angle formed by $[y'_{n-3}, y'_{n-2}]$ and $[y'_{n-3}, B']$ is $\pi/3$. Let y'_{n-1} be the point in the interior of Q such that the length of $[y'_n, y'_{n-1}]$ is equal to $b'/2$ and the angle formed by $[y'_n, y'_{n-1}]$ and $[B', A]'$ is equal to $\pi/3$. Then the length of the horizontal straight segment $[y'_{n-2}, y'_{n-1}]$ is $b'/2$. Now by the choice of the integer n, it follows that the polyline

$$L = [y'_0, \cdots, y'_n]$$

satisfies C_3-bounded linear geometry with $C_3 > 1$ depends only on C_1.

The construction of S' is very similar to that of S. Let $s'_0 = A'$ and $s'_3 = y_{m_{l-1}}$. Let s'_1 be the point in the interior of Q such that the length of $[s'_0, s'_1]$ is equal to $a'/3$ and the angle formed by $[s'_0, s'_1]$ and $[A', B']$ is $\pi/4$. Let s_2 be the point in the interior of Q such that the length of $[s'_2, s'_3]$ is equal to $a'/3$ and the angle formed by $[s'_2, s'_3]$ and $[y'_{m_{l-1}}, A']$ is $\pi/3$. Let S' be the polyline which connects s'_0, s'_1, s'_2 and s'_3.

Let L'_1 bet he part of L' connecting y'_0 and $y'_{m_{l-1}}$ and L'_2 be the remaining part of L'. Then from the construction we see that L' and S' divide Q into four polygons Q_i, $1 \leq i \leq 4$: Q_1 is the top one; Q_2 is the one bounded by L'_2 and $[y_{m_{l-1}}, B']$; Q_3 is the one bounded by L'_1 and S'; and Q_4 is the one bounded by S' and $[A', y_{m_{l-1}}]$. Each of these polygons has four sides, the three of which are straight segments and the last one is a polyline with either C_1 or C_3-bounded linear geometry. Here $C_3 > 1$ is some constant depending only on C_1. Since the geometry of each Q_i is well controlled, it is not difficult to construct the desired qc homeomorphisms ψ_i which maps each Q_i to a standard square with the bottom side being divided into finitely many pieces satisfying either C_1 or C_3-bounded linear geometry.

This proves the Claim and completes the proof of the lemma.

$\square$

Now let us prove Lemma 2.1. The idea of the proof is the same as the one used in [17]. The essential difference is that in the situation here there are two or more critical points in the unit circle, so we will use Lemma 4.13 instead of 4.12. Let $f \in \Pi_{\text{geom}}^{d,\theta_N}$. Let $B \in \mathcal{S}_d$ be the Blaschke product which models f. First let us introduce the concept of Yoccoz's cells. Recall that

$$\mathcal{Q}_n = \{x_i \bigm| 0 \le i < q_n\}.$$

Take

$$N_0 \ge 1 \tag{28}$$

large enough such that for all $n \ge N_0$, $d(x_i, x_j) < 1$ holds for any two adjacent points x_i and x_j in $\mathcal{Q}_n$. Note that N_0 can be taken to be uniform (This is because $\theta_N \to \theta$ and all θ_N and θ are irrational numbers, and $\mathcal{S}_d$ is a compact family). From now on we always assume that $n \ge N_0$.

For each $x_i \in \mathcal{Q}_n$, let y_i be the point on the radial segment $[0, x_i]$ such that

$$|y_i - x_i| = d(x_r, x_l)/2$$

where x_r and x_l denote the two points immediately to the right and left of x_i in $\mathcal{Q}_n$.

Definition 4.2 (Yoccoz's cells)**.** Let x_i and x_j be any two adjacent points in $\mathcal{Q}_n$. Connect y_i and y_j by a straight segment. Then the three straight segments $[x_i, y_i]$, $[y_i, y_j]$, $[x_j, y_j]$, and the arc segment $[x_i, x_j]$ bound a domain in Δ. We call the closure of this domain a *cell of level* n. The segment $[y_i, y_j]$ is called the top side of the cell.

Let E be a cell of level n. Let $E_1, \cdots, E_m$ be all the cells of level $(n+1)$ which are contained in E. Then

$$E \setminus \bigcup_{i=1}^{m} E_i$$

is a polygon. Note that one side of such polygon is a polyline which is the union of the top sides of all the cells of level $(n+1)$ which are contained in E. By Lemmas 4.4 and 4.6 and the construction of the polygons, it follows that the the polyline consists of at most $2d-3$ pieces each of which contains finitely many straight segments satisfying C_1-bounded saddle-node geometry, where $C_1 > 1$ is some uniform constant. Let us call these polygons of type I.

On the other hand, consider the set

$$\mathcal{Q}_n' = \{x_i' \bigm| 0 \le i < q_n\}$$

where $x_i' \in \mathbb{T}$ is the point such that $R_{\theta_N}^i(y_i) = 1$, $0 \le i < q_n$. We can use the same way to define the cells and polygons as above. As above one side of the polygon is a polyline. It is easy to see that the polyline satisfies some C_2-bounded linear geometry with $C_2 > 1$ being some uniform constant. Let us call these polygons of type II.

The construction of the map $H : \Delta \to \Delta$ uses the same idea as in [17]. Suppose P_i is a polygon of type I and level $n \ge N_0$. Let Q_i be the corresponding cell of P_i. We first map P_i and Q_i to polygons described in Lemma 4.13 through qc maps ξ_i and σ_i respectively. Note that the bottoms sides of P_i and Q_i satisfy uniform saddle-node geometry and linear geometry respectively. From Theorem 4.1, ξ_i and σ_i can be constructed so that their qc constants are uniformly bounded (cf. [17]. Since the two bottom sides of $\xi_i(P_i)$ and $\sigma_i(Q_i)$ satisfy uniform saddle and linear geometry, by Lemma 4.13 there is a uniform

$C > 1$ and a qc map $\tau_i : \xi_i(P_i) \to \sigma_i(Q_i)$ such that τ_i maps each edge of $\xi_i(P_i)$ linearly to the corresponding edge of $\sigma_i(Q_i)$ and the qc constant of τ_i is bounded by

$$K_n = C \cdot (1 + (\log a_{n+1})^2).$$

Now define

$$\phi_i = \sigma_i^{-1} \circ \tau_i \circ \xi_i.$$

The complement of all polygons P_i of type I in Δ is a polygon contains the origin in its interior whose boundary is the union of the top sides of all cells of level N_0. Let us denote this polygon by P_0. Similarly, the complement of all polygons Q_i of type II in Δ is a polygon contains the origin in its interior whose boundary is the union of the top sides of all cells of level N_0. Let us denote this polygon by Q_0. Since $\mathcal{S}_d$ is compact, it follows that there is a uniform $K_0 > 1$ such that for all $B \in \mathcal{S}_d$, there is a K_0-qc homeomorphism $\phi_0 : P_0 \to Q_0$ which maps each edge of P_0 linearly to the corresponding edge of Q_0. Now we can define $H : \Delta \to \Delta$ by gluing all ϕ_i along the edges of all polygons P_i.

Let us now prove the existence of the constants $M, \alpha > 0$ and $0 < \epsilon_0 < 1$ so that Lemma 2.1 holds. Let $\epsilon_0 = \frac{2}{1+K_{n_0}}$. By Theorem 4.1, it follows that there exist $C > 1, 0 < \delta < 1$ such that the area of the union of all cells of level n is less that $C_1\delta^n$. For any $0 < \epsilon < \epsilon_0$, let $n \geq N_0$ be the least integer such that $\epsilon > \frac{2}{1+K_{n+1}}$. Then by a simple calculation, it follows that $n > \frac{1}{C_0\epsilon}$ for some uniform constant $C_0 > 0$. By the minimal property of n it follows that $\epsilon \leq \frac{2}{1+K_n}$. This means the following set

$$\{z \in \Delta \mid |\mu_H(z)| > 1 - \epsilon\}$$

is contained in the union of all the cells of level n. Thus the area of the above set is bounded by $C_1\delta^n$. Since $C_1\delta^n = C_1 e^{-n \ln \frac{1}{\delta}} < C_1 e^{-\frac{1}{C_0\epsilon} \cdot \ln \frac{1}{\delta}}$, Lemma 2.1 follows by taking $M = C_1$ and $\alpha = \frac{1}{C_0} \cdot \ln \frac{1}{\delta}$.

4.5. **Proof of Lemma 2.2.** Let $B \in \mathcal{S}_d$ and $h = B|\mathbb{T}$. Let $H : \Delta \to \Delta$ be the David extension of h given in the last subsection. Let

$$\mu_H = \frac{H_{\bar{z}}}{H_z} \frac{d\bar{z}}{dz}$$

be the Beltrami differential given by H. Let μ denote the Beltrami differential on the plane which is the pull back of μ_H by the iteration of $\widehat{B}$.

Let Y_n be the union of all the Yoccoz's cells of level n. Then the outer boundary component of Y_n is $\mathbb{T}$, and the inner boundary component of Y_n is the union of finitely many straight segments, and moreover,

$$Y_{N_0} \supset Y_{N_0+1} \supset \cdots \supset Y_n \supset Y_{n+1} \supset \cdots .$$

From the proof of Lemma 2.1, it follows that

Lemma 4.14. *There exist constants $1 < \lambda < \infty$ such that for any $B \in \mathcal{S}_d$, for all $n \geq N_0$, the dilatation of H in $\Delta \setminus Y_n$ is not greater than $\lambda \cdot n$.*

Define

$$X = \{z \in \mathbb{C} \setminus \overline{\Delta} \bigm| B^k(z) \in \Delta \text{ for some integer } k \geq 1\}.$$

For each $z \in X$, let $k_z \geq 1$ be the least positive integer such that $B^{k_z}(z) \in \Delta$. Define

$$X_n = \{z \in X \bigm| B^{k_z}(z) \in Y_n\}.$$

Lemma 4.15. *There exist* $C_0 > 1$ *and* $0 < \delta_0 < 1$ *such that for any* $B \in \mathcal{S}_d$, $area(X_n) < C_0 \cdot \delta_0^n$ *for all* $n \geq N_0$.

Proposition 4.2. Lemma 4.15 implies Lemma 2.2.

Proof. By Theorem 4.1, there exist uniform constants $C > 0$ and $0 < \delta < 1$ such that

$$area(Y_n) \leq C \cdot \delta^n$$

holds for all $n \geq N_0$. From this and Lemma 4.15 and by taking C and $0 < \delta < 1$ larger we may assume that for all $n \geq N_0$,

$$area(X_n \cup Y_n) < C \cdot \delta^n.$$

Let λ be the constant in Lemma 4.14. Let $\epsilon_0 = \frac{1}{\lambda N_0}$. For any $0 < \epsilon < \epsilon_0$, let $n > N_0$ be the least integer such that $\epsilon > \frac{1}{\lambda(n-1)}$. By Lemma 4.14, if $|\mu(z)| > 1-\epsilon$, then $z \in X_n \cup Y_n$. By the minimal property of n we have $\epsilon \leq \frac{1}{\lambda n}$. It follows that

$$area(\{z \| \mu(z)| > 1-\epsilon\}) < area(X_n \cup Y_n) < C \cdot \delta^n < C \cdot \delta^{\frac{1}{\lambda\epsilon}}$$
$$= C \cdot e^{-(1/\lambda)\cdot(\ln \delta^{-1})/\epsilon}$$

Lemma 2.2 then follows by taking $M = C$ and $\alpha = (1/\lambda) \cdot (\ln \delta^{-1})$. This completes the proof of Proposition 4.2. □

Lemma 4.16. *There exist* $C > 0$, $0 < \epsilon < 1$, $0 < \delta < 1$ *and an integer* $N_1 \geq N_0$ *such that for all* $B \in \mathcal{S}_d$,

$$area(X_{n+2}) \leq C \cdot \epsilon^n + \delta \cdot area(X_n), \quad \forall\, n > N_1. \tag{29}$$

Proposition 4.3. Lemma 4.16 implies Lemma 4.15.

Proof. We may assume that

$$\delta > \epsilon^2$$

since otherwise we may replace δ by $(1+\epsilon^2)/2$ and (29) still holds. Note that every B in $\mathcal{S}_d$ behaves like $z \mapsto z^d$ near infinity. By the compactness of $\mathcal{S}_d$, it follows that there is an $R > 1$ such that $|B(z)| > z$ for all $|z| \geq R$. This implies that X_n is contained in the disk $\{|z| < R\}$ and thus

$$area(X_n) < \pi R^2$$

for all $n \geq 1$. Thus it suffices to prove Lemma 4.15 for $n \geq N_1 + 2$.

From (29) we have

$$area(X_{n+2}) + \frac{C}{\delta - \epsilon^2} \cdot \epsilon^{n+2} \leq \delta \cdot (area(X_n) + \frac{C}{\delta - \epsilon^2} \cdot \epsilon^n).$$

In the case that $n = N_1 + 2k$ for some $k \geq 1$, we have

$$area(X_n) \leq \delta^k (area(X_{N_1}) + \frac{C}{\delta - \epsilon^2} \cdot \epsilon^{N_1}) = (\sqrt{\delta})^{-N_1} \cdot (\pi R^2 + \frac{C}{\delta - \epsilon^2} \cdot \epsilon^{N_1})(\sqrt{\delta})^n.$$

In the case that $n = N_1 + 2k + 1$ for some $k \geq 1$, we have

$$area(X_n) \leq \delta^k (area(X_{N_1+1}) + \frac{C}{\delta - \epsilon^2} \cdot \epsilon^{N_1}) = (\sqrt{\delta})^{(-N_1-1)} \cdot (\pi R^2 + \frac{C}{\delta - \epsilon^2} \cdot \epsilon^{N_1})(\sqrt{\delta})^n.$$

This implies Lemma 4.15. Proposition 4.3 follows. □

The remaining part of this section is devoted to the proof of Lemma 4.16. The basic idea comes from the author's previous work [25]. For $z \in \mathbb{C}$ and $r > 0$, let $B_r(z)$ denote the Euclidean disk with radius r and center z.

Definition 4.3 (K-bounded geometry)**.** *Let $K > 1$ and (U, V) be a pair of sets in $\mathbb{C}$ such that $V \subset U$. We say (U, V) has K-bounded geometry if there exist $x \in V$ and $r > 0$ such that*

$$B_r(x) \subset V \subset U \subset B_{Kr}(x).$$

The following lemma is a variant of Vitali's covering lemma. For a proof, see [25].

Lemma 4.17 (cf. Lemma 2.1 of [25])**.** *Let $K > 1$ and $L = 8K + 9$. Then for any finite family of pairs of measurable sets $\{(U_i, V_i)\}_{i \in \Lambda}$ all of which have K-bounded geometry, namely, for each $i \in \Lambda$, there exist $x_i \in V_i$ and $r_i > 0$ satisfying*

$$B_{r_i}(x_i) \subset V_i \subset U_i \subset B_{Kr_i}(x_i), \tag{30}$$

there is a subfamily σ_0 of Λ such that all $B_{r_j}(x_j), j \in \sigma_0$, are disjoint, and moreover,

$$\bigcup_{i \in \Lambda} U_i \subset \bigcup_{j \in \sigma_0} B_{Lr_j}(x_j).$$

In particular, we have

$$m\Big(\bigcup_{i \in \Lambda} U_i\Big) \leq L^2 \cdot m\Big(\bigcup_{i \in \Lambda} V_i\Big)$$

where $m(\cdot)$ denotes the area with respect to the Euclidean metric.

Recall Δ and $\mathbb{T}$ denote the unit disk and unit circle respectively. Let $\text{diam}(\cdot)$ and $\text{dist}(\ ,\ )$ denote the diameter and distance with respect to the Euclidean metric. Let $\Omega = \mathbb{C} \setminus \overline{\Delta} = \{z \in \mathbb{C} \mid |z| > 1\}$. Then Ω is a hyperbolic Riemann surface.

Definition 4.4. Let $1 < K < \infty$ and $z \in X_{n+2}$. We say z is associated to a K-admissible pair (U, V) if $V \subset U \subset \Omega$ are two open topological disks such that $z \in U$ and

1. $V \subset X_n \setminus X_{n+2}$,
2. the pair (U, V) has K-bounded geometry,
3. $\text{diam}(U) < K \cdot \text{dist}(U, \mathbb{T})$.

Let $I \subset \mathbb{T}$ be an open interval. Let $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$ be the punctured plane. Set

$$\Omega_I = \mathbb{C}^* \setminus (\mathbb{T} \setminus I). \tag{31}$$

Then Ω_I is a hyperbolic Riemann surface. For $d > 0$, the hyperbolic neighborhood of I is defined by

$$\Omega_d(I) = \{z \in \Omega_I \mid d_{\Omega_I}(z, I) < d\} \tag{32}$$

where $d_{\Omega_I}(\ ,\ )$ denotes the hyperbolic distance in Ω_I.

When I is an open interval in the real line $\mathbb{R}$, the hyperbolic neighborhood of I in the slit plane $\mathbb{C} \setminus (\mathbb{R} \setminus I)$ is bounded by two arc segments of Euclidean circles which are symmetric about $\mathbb{R}$, and moreover, the exterior angles formed by the arcs and $\mathbb{R}$ are all equal and determined by the formula $d = \log \cot(\alpha/4)$. The next lemma says when I is small enough, $\Omega_d(I)$ is very much like the hyperbolic neighborhood in the slit plane.

Lemma 4.18 (cf. Lemma 2.2 of [25])**.** *Let $d > 0$ be given. Then when I is small, $\Omega_d(I)$ is a Jordan domain and is like the hyperbolic neighborhood of the slit plane in the following sense.*

1. $\partial\Omega_d(I) = \gamma_{int} \cup \gamma_{out}$ *where γ_{int} and γ_{out} are two real analytic curve segments both of which connect the two end points of I. Moreover, $\gamma_{int} \setminus \partial I \subset \Delta$ and $\gamma_{out} \setminus \partial I \subset \mathbb{C} \setminus \overline{\Delta}$;*
2. *γ_{int} and γ_{out} are symmetric about $\mathbb{T}$, and each of them is like an arc segment of some Euclidean circle;*
3. *Let α denote the exterior angles formed by γ_{int} and $\mathbb{T}$, γ_{out} and $\mathbb{T}$, all of which are the same, then $d = \log\cot(\alpha/4)$.*

Note that $\Omega_d(I)$ is divided by I into two parts: one is in the interior of Δ and the other one is in the exterior of Δ. We only consider the part which is in the exterior of Δ. Let $H_\alpha(I)$ denote this part. That is,

$$H_\alpha(I) = \{z \in \Omega_d(I) \mid |z| > 1\} \tag{33}$$

where α is determined by the formula $d = \log\cot(\alpha/4)$.

Let

$$\Pi_n(B) = \{I_n^i,\ 0 \le i \le q_{n+1} - 1;\ \ I_{n+1}^i,\ 0 \le i \le q_n - 1\}$$

be the collection of intervals defined in (22). Define

$$Z_n = \bigcup_{0 \le i \le q_{n+1}-1} H_\alpha(I_n^i) \cup \bigcup_{0 \le i \le q_n - 1} H_\alpha(I_{n+1}^i). \tag{34}$$

It is easy to see that Z_n is the outer half of an open neighborhood of $\mathbb{T}$. See Figure 2 for an illustration. As a consequence of Theorem 4.1 and the fact that $\mathrm{diam}(H_\alpha(I)) = O(|I|)$ (cf. Lemma 4.18) , it follows that there exist $C > 1$ and $0 < \epsilon < 1$ depending only on d such that for all $B \in \mathcal{S}_d$ and $n \ge 1$,

$$area(Z_n) < C \cdot \epsilon^n. \tag{35}$$

Let us first reduce Lemma 4.16 to

Lemma 4.19. *There exist $K > 1$ and $N_1 \ge N_0$ such that for all $B \in \mathcal{S}_d$ and $n \ge N_1$, if $z \in X_{n+2}$, then either $z \in Z_n$, or z is associated to some K-admissible pair (U, V).*

Proposition 4.4. Lemma 4.19 implies Lemma 4.16.

Proof. From the proof of Proposition 4.3 we have an $R > 1$ such that for all $n \ge 1$ we have

$$X_n \subset \{z \mid |z| \le R\}.$$

Note that X_{n+2} is a Lebesgue measurable set. So for any $\eta > 0$, by a standard result in Lebesgue measure theory, there is a closed set $F \subset X_{n+2}$ such that

$$area(X_{n+2} \setminus F) < \eta. \tag{36}$$

Since Z_n is open and F is a bounded and closed set, it follows that $F \setminus Z_n$ is a compact set.

For any point $x \in F \setminus Z_n$, by Lemma 4.19, x is associated to some K-admissible pair (U_i, V_i) for some uniform $1 < K < \infty$. This implies that the sets U_is form an open cover



FIGURE 2. The set Z_n

of $F \setminus Z_n$. Since $F \setminus Z_n$ is a compact set, we have finitely many pairs (U_i, V_i), $i \in \Lambda$, such that

$$F \setminus Z_n \subset \bigcup_{i \in \Lambda} U_i. \tag{37}$$

From (36) and (37), we have

$$area(X_{n+2}) \le area(Z_n) + area(\bigcup_{i \in \Lambda} U_i) + \eta. \tag{38}$$

By Lemma 4.17, we have

$$area(\bigcup_{i \in \Lambda} U_i) \le \lambda(K) \cdot area(\bigcup_{i \in \Lambda} V_i) \tag{39}$$

where $\lambda(K) = (8K+9)^2$. Since $V_i \subset X_n \setminus X_{n+2}$ for all $i \in \Lambda$, we have

$$area(\bigcup_{i \in \Lambda} V_i) \le area(X_n) - area(X_{n+2}).$$

This, together with (39), implies that

$$area(\bigcup_{i \in \Lambda} U_i) \le \lambda(K) \cdot (area(X_n) - area(X_{n+2})). \tag{40}$$

From (38) and (40) we get

$$(1 + \lambda(K)) \cdot area(X_{n+2}) \le area(Z_n) + \lambda \cdot area(X_n) + \eta.$$

From (35) and the above inequality we have

$$area(X_{n+2}) \le \frac{\lambda(K)}{1+\lambda(K)} \cdot area(X_n) + \frac{C}{1+\lambda(K)} \cdot \epsilon^n + \frac{\eta}{1+\lambda(K)}. \tag{41}$$

Since $\eta > 0$ is arbitrary, by letting $\eta \to 0$, we get

$$area(X_{n+2}) \le \frac{\lambda(K)}{1+\lambda(K)} \cdot area(X_n) + \frac{C}{1+\lambda(K)} \cdot \epsilon^n.$$

The implies Lemma 4.16. The proof of Proposition 4.4 is completed.

$\square$

Lemma 4.20. *Let $1 < L < \infty$. Then there is a $1 < K < \infty$ depending only on L such that for any $B \in \mathcal{S}_d$, $z \in X_{n+2}$ and $m \ge 1$, if $B^i(z) \in \mathbb{C} \setminus \overline{\Delta}$ for all $1 \le i \le m$ and $\zeta = B^m(z)$ is associated to some L-admissible pair (U, V), then z is associated to some K-admissible pair.*

Proof. This is a direct consequence of the Koebe's distortion theorem. $\square$

Let us now begin the proof of Lemma 4.19. Let $B \in \mathcal{S}_d$ and $z \in X_{n+2}$. Suppose $z \notin Z_n$. It suffices to prove that there is a $1 < K < \infty$ independent of B, n and z such that z is associated to some K-admissible pair (U, V). Recall that $k_z \ge 1$ is the least positive integer such that $B^{k_z}(z) \in \Delta$. Since $z \in X_{n+2}$ we have $B^{k_z}(z) \in Y_{n+2}$. Let us denote

$$z_l = B^l(z), \; 0 \le l \le k_z.$$

Since $z_0 = z \notin Z_n$, the set

$$\Pi = \{k \in \mathbb{Z} \mid 0 \le k < k_z \text{ and } B^k(z) \notin Z_n\}$$

is not empty. It is clear that Π contains at most k_z elements and is thus a finite set. Let

$$k_0 = \max_{k \in \Pi}\{k\}.$$

Then $0 \le k_0 \le k_z - 1$. Set

$$\zeta = z_{k_0} \text{ and } \omega = B(\zeta) = z_{k_0+1}.$$

Then $\zeta \notin Z_n$, and moreover,

$$\omega \in Z_n \text{ if } k_0 < k_z - 1 \text{ and } \omega \in Y_{n+2} \text{ if } k_0 = k_z - 1. \tag{42}$$

By Lemma 4.20 it suffices to prove that ζ is associated to some L-admissible pair (U, V) for some uniform $1 < L < \infty$ depending only on d. The proof is divided into two cases.

Note that there are exactly $d-1$ Jordan curves in the outside of Δ, each of which is a pre-image of $\mathbb{T}$ and is attached to $\mathbb{T}$ at a critical point c. Note that when n is large, ω is close to $\mathbb{T}$ uniformly. There are $d-1$ pre-images of ω close to the $d-1$ Jordan curves respectively, and the last one is close to $\mathbb{T}$. In the case that $\omega \in Z_n$, the last one lies in the exterior of Δ, and in the case that $\omega \in Y_{n+2}$, the last one lies in Δ.

Let I be the arc in the dynamical partition of level n such that $\omega \in H_\alpha(I)$ or ω belongs to the cell E which is attached to I or the union of I and one of its adjacent intervals in the dynamical partition of level n. Let us first group the critical points such that the ones which are closed to each other are classified into one group. Since there are at most $d-1$ critical values of B in $\mathbb{T}$, for any $L > 1$, we have $0 < \eta(L, d) < 1$ such that for any B, we can find $\eta(L, d) < \epsilon < \delta < 1$ with $\delta/\epsilon > L$ and for each critical value v of B in $\mathbb{T}$, we either have $\text{dist}(\omega, v) < \epsilon \cdot |I|$ or $\text{dist}(\omega, v) > \delta \cdot |I|$. We have two cases.

Case I. There is a critical value, say v, such that $\text{dist}(\omega, v) < \epsilon \cdot |I|$. By the choice of ϵ and δ, for each critical value v of B in $\mathbb{T}$, we either have $\text{dist}(\omega, v) < \epsilon \cdot |I|$ or

$\operatorname{dist}(\omega, v) > \delta \cdot |I|$. By the geometry of the cells (cf. Lemma 6.3 of [17] or Lemma 6.1 of [25]), we have a pair of Jordan domains $B \subset A$ such that the following properties hold.

1. $B \subset E \setminus Y_{n+2}$ is a Euclidean disk with radius $\asymp |I|$,
2. A contains ω and all v with $\operatorname{dist}(\omega, v) < \epsilon \cdot |I|$,
3. for all other v, $\operatorname{dist}(v, A) \asymp \operatorname{diam}(A) \asymp |I|$ and for any point $z \in A$, the angle formed by $[v, z]$ and $\mathbb{T}$ is less than $\kappa \cdot \pi$ with $0 < \kappa < 1$ being some constant depending only on $\mathcal{S}_d$.

For critical point c corresponding to critical value v bounded away from A, the preimage of A which intersects the corresponding pre-circle at c is a desired pair by Koebe's distortion theorem. In fact, let $B' \subset A'$ be the preimages of $B \subset A$ respectively. Then A' has bounded hyperbolic diameter. Thus $B' \subset A'$ has K-bounded geometry for some uniform $K > 1$. If ζ belongs to these A', the lemma has been proved. Otherwise, let us prove ζ belongs to Z_n. To see this, use a simple closed geodesic γ in $\widehat{\mathbb{C}} \setminus V'$, with V' being the union of the set of critical values and the two end points of I, turning around ω and all those critical values with $\operatorname{dist}(\omega, v) < \epsilon \cdot |I|$, and such that at least one of the end points of I lies in the outside of γ (This requirement implies that γ intersects I). Because $\delta/\epsilon > L$, by taking L large, we can make sure there is such a γ which can be arbitrarily short provided that L is large enough. Let $J \subset \mathbb{T}$ be the arc such that $B(J) = I$. Let η be the pre-image of γ which intersects J. Then η is a simple closed geodesic in $\widehat{\mathbb{C}} \setminus B^{-1}(V')$ and can be arbitrarily short provided that L is large enough. In particular, compared with $|J|$, the Euclidean diameter of η can be arbitrarily small provided that L is large enough. By consider the covering degree of $B : \eta \to \gamma$, it follows that all the other pre-images of ω are contained in the inside of η. By the construction of Z_n, it follows that Z_n contains a definite neighborhood of J. Thus Z_n contains η. This implies that all the other η is pre-images of ω belongs to Z_n. This proves the lemma in the first case.

Case II. $\operatorname{dist}(\omega, v) > \delta|I|$ for all critical values v.

Subcase I of Case II: I contains no critical values and $I \neq I_{n+1}^{q_n - 1}$. Let $J \subset \mathbb{T}$ be the arc such that $B(J) = I$. Since I does not contain $B(1)$, $I \neq I_n^{q_{n+1}-1}$. So J is one of the intervals in the dynamical partition of level n. We can construct a pair of Euclidean disk $B \subset A$ with $\omega \in A$ such that $B \subset E \setminus Y_{n+2}$ and

1. $A \setminus \overline{\Delta} \subset H_\alpha(I)$,
2. $\operatorname{dist}(v, A) \succeq \operatorname{diam}(A) \asymp \operatorname{diam}(B) \asymp |I|$,
3. A has a Jordan neighborhood $\tilde{A}$ which contains no critical values and such that $\tilde{A} \cap \mathbb{T}$ is connected arc segment and $\tilde{A} \setminus \overline{A}$ has a definite positive modulus.

There are two subcases. Let U be the component of $B^{-1}(A)$ which intersects J. Since $A \setminus \overline{\Delta} \subset H_\alpha(I)$, by Schwarz lemma, $U \setminus \overline{\Delta} \subset H_\alpha(J) \subset Z_n$. This implies the pre-image of ω contained in U is contained in Z_n. Note that there are exactly $d-1$ other components of $B^{-1}(A)$ each of which intersects a pre-circle at some critical point $c \in \mathbb{T}$. Suppose $\zeta \notin U$ since otherwise we have proved the lemma. Then ζ must belong to one of these $d-1$ components, say A'. Let $B' \subset A'$ be the pre-image of B contained in A'. Let us explain that (A', B') is a desired pair. To see this, note that the inverse branch of B can be holomorphically extended to $\tilde{A}$. Since $\tilde{A} \setminus \overline{A}$ has a definite positive modulus, thus A' has a protective collar in the exterior of Δ with a definite positive modulus. Since

$\mathrm{diam}(B) \asymp \mathrm{diam}(A)$, it follows from Koebe's distortion theorem that B' contains a disk whose diameter is uniformly commensurable with that of A'. Since A' has a protective collar in the exterior of Δ which has definite modulus, we have $\mathrm{dist}(A', \mathbb{T}) \succeq \mathrm{diam}(A')$.

Subcase II of Case II: $\mathrm{dist}(\omega, \mathbb{T}) < \epsilon|I|$. Let ω' be the point in $\mathbb{T}$ which is nearest to ω. Let X be the set consisting of all critical values, the two end points of I and ω, ω'. Since $\mathrm{dist}(\omega, \omega') < \epsilon|I|$, we have a short simple closed geodesic γ in $\widehat{\mathbb{C}} \setminus X$ which intersects I and enclose ω and ω' in its inside. The pre-image of γ which intersects J, is a short simple closed geodesic in $\widehat{\mathbb{C}} \setminus B^{-1}(X)$. Note that η encloses a pre-image of ω, say ζ_0. Since η is short, it follows that such η can be contained in an arbitrarily small neighborhood of J provided that ϵ is small enough. Since Z_n contains a definite neighborhood of J, it follows that Z_n contains ζ_0. Let us assume that $\zeta \neq \zeta_0$, since otherwise the lemma has been proved. Let us consider the remaining $(d-1)$ pre-images of ω. Since $\mathrm{dist}(\omega, v) > \delta|I|$ and $\mathrm{dist}(\omega, \mathbb{T}) < \epsilon|I|$, we can still construct a Euclidean disk $B \subset E \setminus Y_{n+2}$ and a Jordan domain A containing ω and ω' such that $B \subset A$ and the requirements (1), (2), and (3) in the proof of Subcase I of Case II are satisfied. There are $(d-1)$ pre-images of ω which are contained in the $(d-1)$ components of $B^{-1}(A)$ intersecting the $(d-1)$ precircles at the $(d-1)$ critical points in $\mathbb{T}$. As in the proof of the Subcase I of Case II, we have seen if ζ is one of these pre-images of ω, then ζ is associated to some K-admissible pair. This proves the lemma in the Subcase II of Case II.

Subcase III of Case II: $\mathrm{dist}(\omega, \mathbb{T}) \ge \epsilon|I|$ and I contains at least one critical value or $I = I_{n+1}^{q_n-1}$. So either I or one of its adjacent intervals in the dynamical partition of level n contains critical value(s) of B. Then there exists a disk $B \subset E \setminus Y_{n+2}$ and two Jordan domains, say A_0 and A_1, which contain ω and B such that

1. $\mathrm{dist}(A_i, v) \asymp \mathrm{diam}(A_i) \asymp |I|$ for $i = 0, 1$ and every critical value $v \in \mathbb{T}$,
2. $A_0 \cup A_1$ separates at least one critical value from ∞ (That is, A_0 and A_1 turn around some nearby critical value(s) in an opposite way),
3. For $i = 0, 1$, A_i has a Jordan neighborhood $\tilde{A}_i$ which contains no critical values and such that $\tilde{A}_i \cap \mathbb{T}$ is connected arc segment and $\tilde{A}_i \setminus \overline{A_i}$ has a definite positive modulus.

Here we would like to emphasize that the condition $\mathrm{dist}(\omega, \mathbb{T}) \ge \epsilon|I|$ is used to meet the requirement (3). This is because if ω is too close to $\mathbb{T}$, then one of $\tilde{A}_i \cap \mathbb{T}$, $i = 1, 2$ must contain at least two intervals which are separated by some critical value and thus can not be a connected arc segment. Now first note that there are $d-1$ components of $B^{-1}(A_0)$ each of which intersects a precircle at a critical point at $\mathbb{T}$ and contains a pre-image of ω. If ζ belong to one of these components, as we have seen in the proof of Subcase I of Case II, ζ is associated to some K-admissible pair (U, V). Since there are d pre-images of ω, it suffices to deal with the last pre-image of ω. To do this, let $v_1, \cdots, v_m$ be all the critical values which are separated by $A_0 \cup A_1$ from ∞. Assume that v_1 is the one which is closest to $A_0 \cap \mathbb{T}$. Let c_1 be the critical point such that $B(c_1) = v_1$. Let A_1' be the component of $B^{-1}(A_1)$ which intersects the pre-circle at c_1. Then A_1' contains a pre-image of ω. It is important to note that this pre-image is different from the previous $(d-1)$ preimages: this is because for each of the previous $(d-1)$ pre-images of ω, the associated inverse branch of B maps $A_0 \cap \mathbb{T}$ to a curve segment on one of the $(d-1)$ pre-circle at sone critical point in $\mathbb{T}$, and for the last pre-image, the inverse branch of B

defined on A_1 can be extended to $A_0 \cap \mathbb{T}$ and maps it to $\mathbb{T}$. Let B_1' be the pre-image of B which is contained in A_1'. Then (A_1', B_1') is the desired pair to which the last pre-image of ω is associated. This proved the lemma in the Subcase III of Case II.

This completes the proof of Lemma 4.19. Lemma 2.2 thus follows.

4.6. **Proof of the Key-Lemma 1.** Note that $H_B^{-1}(0)$ is uniformly bounded away from the unit circle. Let ϕ be the David homeomorphism mapping 0 to $H^{-1}(0)$ and fix 1 and ∞. Using Petersen-Zakeri's argument as in the proof of Lemma 5.5 of [17], it follows that $\phi \circ \widehat{B} \circ \phi^{-1}$ is holomorphic and thus belong to $\Pi_{geom}^{\alpha,d}$. By the rigidity result in §6, every map in this class can be obtained in this way. By Lemmas 2.2 and 2.3 it follows that ϕ belongs to a normal family. The existence of λ and η can be proved by a compactness argument. For $0 < \delta', \delta < 2$, define

$$\lambda(\delta) = \sup |P^k(1) - P^m(1)|$$

where sup is taken over all $B \in \mathcal{S}_d$ and all $k > m \geq 0$ such that $|e^{2\pi i k\alpha} - e^{2m\pi i\alpha}| < \delta$, and

$$\eta(\delta') = \inf |P^k(1) - P^m(1)|$$

where inf is taken over all $B \in \mathcal{S}_d$ and all $k > m \geq 0$ such that $|e^{2\pi i k\alpha} - e^{2m\pi i\alpha}| > \delta'$. It is sufficient to prove $\eta(\delta') > 0$ and $lim \lambda(\delta) \to 0$ as $\delta \to 0$. Suppose $\eta(\delta') = 0$ for some $\delta' > 0$. Then there is a sequence $h_n \to h$, and $\phi_n \to \phi$ and k_n, m_n. This implies a contradiction. The same argument implies that $\eta(\delta') > 0$. The proof of Key-Lemma 1 is completed.

5. Proofs of the Key-Lemmas 3 and 4

Let $m, l \geq 1$ be two integers and

$$\zeta_0 = (w_1^0, \cdots, w_m^0, w_{m+1}^0, \cdots, w_{m+l}^0)$$

be a point in $\mathbb{C}^{m+l}$. Let $\xi_0 = (w_{m+1}^0, \cdots, w_{m+l}^0)$ be the point in $\mathbb{C}^l$. Let

$$Q_i(w_1, \cdots, w_m, w_{m+1}, \cdots, w_{m+l}), \ \ 1 \leq i \leq m,$$

be polynomials of $m + l$ complex variables.

Suppose that there exist an open neighborhood U of ξ_0 in $\mathbb{C}^l$ and m functions

$$w_i = g_i(w_{m+1}, \cdots, w_{m+l}), \ \ 1 \leq i \leq m,$$

defined in U such that for all $(w_{m+1}, \cdots, w_{m+l}) \in U$, we have

$$Q_i(g_1, \cdots, g_m, w_{m+1}, \cdots, w_{m+l}) = 0,$$

and moreover, there is an open neighborhood V of $(w_1^0, \cdots, w_m^0)$ in $\mathbb{C}^m$ such that for any points $(w_{m+1}', \cdots, w_{m+l}') \in U$ and $(w_1', \cdots, w_m') \in V$, if

$$Q_i(w_1', \cdots, w_m', w_{m+1}', \cdots, w_{m+l}') = 0, \ \ 1 \leq i \leq m,$$

then

$$w_i' = g_i(w_{m+1}', \cdots, w_{m+l}'), \ \ 1 \leq i \leq m.$$

We first prove such functions g_i are determined by polynomial equations. More precisely,

Lemma 5.1. *There exist m irreducible polynomials P_i, $1 \le i \le m$, of $m+1$ variables such that*

$$P_i(g_i, w_{m+1}, \cdots, w_{m+l}) = 0, \;\; 1 \le i \le m.$$

Proof. Let us start from the system of polynomial equations

$$Q_i(w_1, \cdots, w_m, w_{m+1}, \cdots, w_{m+l}) = 0, \; 1 \le i \le m. \tag{43}$$

By the definition of g_i, it follows that

$$Q_i(g_1, \cdots, g_m, w_{m+1}, \cdots, w_{m+l}) = 0$$

for all $(w_{m+1}, \cdots, w_{m+l}) \in U$. So if some Q_i is not irreducible, then Q_i has an irreducible factor, say S, such that $S(g_1, \cdots, g_m, w_{m+1}, \cdots, w_{m+l}) = 0$ for all $(w_{m+1}, \cdots, w_{m+l}) \in U$. Then in the system (43), we replace Q_i by S. In this way we may assume that all Q_i are irreducible.

Let us first write all Q_i into polynomials of w_1 with coefficients being polynomials of $w_2, \cdots, w_{m+l}$. If some Q_i has a term with coefficient, say

$$h(w_2, \cdots, w_m, w_{m+1}, w_{m+l}),$$

satisfying

$$h(g_2, \cdots, g_m, w_{m+1}, \cdots, w_{m+l}), w_{m+1}, \cdots, w_{m+l}) = 0$$

for all $(w_{m+1}, \cdots, w_{m+l}) \in U$, then we add the equation

$$h(w_2, \cdots, w_{m+l}) = 0$$

to the system (43), and at the same time, delete the corresponding term from Q_i. Repeating the argument in the beginning of the proof if necessary, we can make sure all the polynomials in the new system are still irreducible. Besides this, if one polynomial is the constant multiple of the other, we just remove one of them from the system. In this way we always make sure the system is not redundant.

We claim that there is at least one polynomial in the new system which involves w_1. Since otherwise the polynomial equations in the new system will hold in U by taking

$$w_i = g_i(w_{m+1}, \cdots, w_{m+l}), \;\; 2 \le i \le m,$$

and taking w_1 to be any function of $(w_{m+1}, \cdots, w_{m+l})$. This then implies the origin system (43) holds under the same assignment of w_i, $1 \le i \le m$. But this contradicts with the uniqueness part in the Implicit Function Theorem. The claim has been proved.

Now let Q_1 be a polynomial in the new system which involves w_1 and moreover, the degree of Q_1 with respect to w_1 is the lowest. Assume there is some other polynomial in the system which also involves w_1. Otherwise we go to the next step. For any other polynomial which involves w_1, say Q_2, we have

$$Q_2 = D \cdot Q_1 + R \tag{44}$$

where D and R are polynomials of w_1 with coefficients being rational functions of $w_2, \cdots, w_{m+l}$, and moreover, the degree of R with respect to w_1 is less than that of Q_1. Since all Q_i are irreducible and the system is not redundant, R is not identically zero. From the above process, we also know if $D = D'/D''$ and $R = R'/R''$ then

$$D''(g_2, \cdots, g_m, w_{m+1}, \cdots, w_{m+l}) \text{ and } R''(g_2, \cdots, g_m, w_{m+1}, \cdots, w_{m+l})$$

is not identically zero in U. Now we replace Q_2 by R'. Again repeat the procedure we previously used, we may assume that the polynomials in the new system are all irreducible and are not redundant. From (44) it follows that under the condition that $R'' \neq 0$ and $D'' \neq 0$, the two equations $R' = 0$ and $Q_1 = 0$ imply $Q_2 = 0$. Since R'' and D'' are not identically zero in U, if $\tilde{g}_i$, $1 \leq i \leq m$, are a group of solutions of the new system and close enough to g_i, $1 \leq i \leq m$, such that R'' and D'' do not vanish in some open subset of U, then they are also a group of solutions of the original system in the open subset of U.

Note that after the above process, the total degree of w_1 is decreased at least by 1. So after finitely many steps, we have a system of irreducible polynomial equations such that

1. there is only one polynomial which involves w_1,
2. $w_i = g_i$, $1 \leq i \leq m$, are a group of solutions of the system,
3. if $w_i = \tilde{g}_i$, $1 \leq i \leq m$, are a group of solutions of the system defined in some open subset in U, which are close enough to g_i, then $\tilde{g}_i = g_i$ for all $1 \leq i \leq m$.

We claim, except Q_1, there is at least one polynomial in the system which involves w_2. Since otherwise, we may let $\tilde{g}_i = g_i$ for $3 \leq i \leq m$, and let $\tilde{g}_2$ be any holomorphic function in U which is close to g_2 but not identical with g_2, then solve $\tilde{g}_1$ by $Q_1(\tilde{g}_1, \tilde{g}_2, g_3, \cdots, g_m, w_{m+1}, \cdots, w_{m+l}) = 0$ and get a solution $\tilde{g}_1$ in some open subset of U which can be arbitrarily close to g_1 provided that $\tilde{g}_2$ is close enough to g_2. So

$$R''(\tilde{g}_1, \cdots, \tilde{g}_m, w_m, \cdots, w_{m+l}) \text{ and } D''(\tilde{g}_1, \cdots, \tilde{g}_m, w_m, \cdots, w_{m+l})$$

do not vanish in this open subset of U. By the third property above we get $\tilde{g}_2 = g_2$. This is a contradiction.

Now consider all the polynomial equations except Q_1 in the system. Let Q_2 be a polynomial in the system which involves w_2 and whose degree with respect to w_2 is the lowest. For each other polynomial except Q_1 which involves w_2, we repeat the previous procedure. Then again after finitely many times we get a new system such that

1. except Q_1, there is only one polynomial Q_2 which involves w_2,
2. $w_i = g_i$, $1 \leq i \leq m$, are a group of solutions of the system,
3. if $w_i = \tilde{g}_i$, $1 \leq i \leq m$, are a group of solutions of the system defined in some open subset in U, which are close enough to g_i, then $\tilde{g}_i = g_i$ for all $1 \leq i \leq m$.

Repeating this procedure finitely many times, we finally get a system which consists of exactly m irreducible polynomial equations, say

$$\begin{gathered} Q_1(w_1, \cdots, w_m, w_{m+1}, \cdots, w_{m+l}) = 0, \\ Q_2(w_2, \cdots, w_m, w_{m+1}, \cdots, w_{m+l}) = 0, \\ \cdots \\ Q_m(w_m, w_{m+1}, \cdots, w_{m+l}) = 0. \end{gathered}$$

Let $P_m = Q_m$. Then $w_m = g_m$ satisfies an irreducible polynomial equation

$$P_m(g_m, w_{m+1}, \cdots, w_{m+l}) = 0.$$

By relabeling each of $w_1, \cdots, w_{m-1}$ as w_m and repeating the above process, we get an irreducible polynomial P_i of $m+1$ variables such that

$$P_i(g_i, w_{m+1}, \cdots, w_{m+l}) = 0, \ \ 1 \leq i \leq m-1.$$

The proof of Lemma 5.1 is completed. □

5.1. **Proof of the Key-Lemma 3.** Recall that $c_{d-1} = 1$. Let x_i be one of the points in the i-th periodic attracting cycle and $p_i \geq 1$ be the period. Since x_i is a periodic point of f with period p_i with multiplier t_i, we have

$$f^{p_i}_{c_1^0,\cdots,c_{d-2}^0,1}(x_i) - x_i = 0 \text{ and } Df^{p_i}_{c_1^0,\cdots,c_{d-2}^0,1}(x_i) - t_i = 0.$$

Since the cycle containing x_i is attracting, there exist open neighborhoods U of $(c_1^0, \cdots, c_{d-2}^0)$ and open neighborhood V of (x_i, t_i) and two holomorphic functions

$$x = g_1(c_1, \cdots, c_{d-2}) \text{ and } t = g_2(c_1, \cdots, c_{d-2})$$

defined in U such that for all $(c_1, \cdots, c_{d-2}) \in U$ we have

$$f^{p_i}_{c_1,\cdots,c_{d-2},1}(x) - x = 0 \text{ and } Df^{p_i}_{c_1,\cdots,c_{d-2},1}(x) - t = 0,$$

and moreover, for any $(C_1', \cdots, c_{d-2}') \in U$ and $(x', t') \in V$ satisfying the above two equations, we have

$$x' = g_1(c_1', \cdots, c_{d-2}') \text{ and } t' = g_2(c_1', \cdots, c_{d-2}').$$

Now apply Lemma 5.1, we have an irreducible polynomial of $d-1$ variables, say Q_i, such that the function $t = g_2(c_1, \cdots, c_{d-2})$ is determined by the following polynomial equation:

$$Q_i(t_i, c_1, \cdots, c_{d-2}) = 0.$$

The map

$$\Phi : (c_1, \cdots, c_l, c_{l+1}, \cdots, c_{d-2}) \to (t_1, \cdots, t_l)$$

is holomorphic. On the other hand, we can construct a holomorphic map

$$\Psi : (t_1, \cdots, t_l) \to (c_1, \cdots, c_{d-1})$$

such that the composition

$$\Phi \circ \Psi = id.$$

The construction can be done as following. Take a point x_i in each cycle. Assume that $t_i \neq 0$ first. Take a small neighborhood of x_i such that $f^{p_i} : U \to U$ is conjugate to the linear map $z \to t_i z$, and moreover, ∂U is an analytic curve. Let $\Gamma = \partial U$ and $\Gamma' = f^{p_i}(\Gamma)$. Then Γ and Γ' bound an annulus. In a small open open disk centered at t_i, define $f_t = f$ on Γ and f_t be a natural holomorphic motion of Γ' as t varies in the disk. Such holomorphic motion can be extended to the interior of the annulus. Here natural means it maps x_i to $f(x_i)$ but with multiplier different, and all the others are the same. In this way we construct ϕ_t depends holomorphically onto t. The map $\phi^{-1} \circ f_t \circ \phi_t$ is the desired map.

This means the Jacobi matrix of $(t_1, \cdots, t_l)$ to $(c_1, \cdots, c_l)$ are non-degenerate by relabeling c_1, c_{d-1} if necessary. This implies that there exist holomorphic functions g_i, $1 \leq i \leq l$, such that

$$Q_i(t_i, g_1, \cdots, g_l, c_{l+1}, \cdots, c_{d-2}) = 0.$$

Moreover, these functions has the desired uniqueness property in a small neighborhood. Applying Lemma 5.1, the proof of the Key-Lemma 3 is completed.

5.2. **Proof of the Key-Lemma 4.** Choose one of the attracting periodic cycles. As in the end of the proof of the Key-Lemma 3, we can construct a holomorphic family of polynomials f_t of one parameter which preserves all the orbit relations and the multipliers of all other attracting cycles. In this way all c_i, $1 \le i \le d-2$ are holomorphic functions of t when t varies in a small neighborhood of t_i. Take one of $c_i(t)$ which is not a constant function. Then there exists a small open disk which can be arbitrarily close to t_i, the function $t \to c_i(t)$ is one-to-one. This implies that in this small disk, $t = \phi(c_i)$ where ϕ is some holomorphic function of c_i. Thus all other c_j are functions of c_i. Thus we have $P(c_j, c_i) = 0$ for some polynomial P. This proves the second implicit theorem.

6. TOPOLOGICAL CHARACTERIZATION OF THE MAPS IN $\Sigma_{geom}^{\alpha,d}$

As we have seen in the previous sections, the maps in the class $\Sigma_{geom}^{\alpha,d}$ play a fundamental role in our proof. The purpose of this section is to establish the characterization and the rigidity of the maps in this class. As a consequence we will prove the Key-Lemma 2. One of the main tools used in this section is a characterization theorem for sub-hyperbolic rational maps obtained in [4] and [26]. This result is an extension of Thurston's characterization of post-critically finite rational maps.

6.1. **Characterization of sub-hyperbolic rational maps.** Let $F : \widehat{\mathbb{C}} \to \widehat{\mathbb{C}}$ be a finitely branched covering map which preserves the orientation. Let

$$\Omega_F = \{z \in \widehat{\mathbb{C}} \mid \deg_z(f) \ge 2\}$$

and

$$P_F = \overline{\bigcup_{k \ge 1} F^k(\Omega_F)}$$

be the critical set and the post-critical set of F. We say F is geometrically finite if P_f is an infinite set but the accumulation set of P_F is a finite set. It is easy to check that each accumulation point of P_F is a period point of F. Let F be a geometrically finite branched covering map. We say F is semi-rational if for each periodic cycle $\mathcal{O} \subset P_F$, there is an open neighborhood U of $\mathcal{O}$ such that $F|U$ is holomorphic, and moreover, $|DF^p(x)| < 1$ for each $x \in \mathcal{O}$ where $p \ge 1$ is the period of $\mathcal{O}$.

Let F be a geometrically finite and semi-rational branched covering map. Let $\gamma \subset \widehat{\mathbb{C}} \setminus P_F$ be a simple closed curve. We say γ is non-peripheral if each component of $\widehat{\mathbb{C}} \setminus \gamma$ contains at least two points in P_F. A multi-curve Γ is a finite family of non-peripheral curves which are disjoint and non-homotopic to each other. We say Γ is stable if for each $\gamma_i \in \Gamma$, each non-peripheral component of $F^{-1}(\gamma)$ is homotopic to some γ_j in Γ.

Suppose $\Gamma = \{\gamma_1, \cdots, \gamma_n\}$ is a stable multi-curve. For $1 \le i, j \le n$, let $\gamma_{i,j,\alpha}$ denote all the non-peripheral components of $f^{-1}(\gamma_i)$ which are homotopic to γ_j. Let $d_{i,j,\alpha}$ be the covering degree of

$$F : \gamma_{i,j,\alpha} \to \gamma_i.$$

Let

$$a_{i,j} = \sum_{\alpha} \frac{1}{d_{i,j,\alpha}}.$$

The matrix $A = (a_{i,j})$ is called Thurston linear transformation matrix. It is non-negative and thus has a maximal positive eigenvalue $\lambda > 0$. If $\lambda \geq 1$ we call Γ a Thurston obstruction.

Definition 6.1. *Two semi-branched and geometrically finite branched covering maps F and G are called CLH-equivalent (combinatorially and locally holomorphically equivalent) if there exist a pair of homeomorphisms of the sphere $\phi, \psi : \widehat{\mathbb{C}} \to \widehat{\mathbb{C}}$ such that*

1. $\phi \circ F = G \circ \psi$,
2. *For each holomorphic attracting cycle $\mathcal{O}$, there is an open neighborhood U of $\mathcal{O}$, say U, such that $\phi|U = \psi|U$ are holomorphic,*
3. *ϕ is isotopic to ϕ rel $P_F \cup \cup_i \overline{U_i}$.*

The following theorem is an extension of Thurston's characterization for post-critically finite rational maps to sub-hyperbolic rational maps.

Theorem 6.1 ([4] & [26])**.** *Let F be a geometrically finite and semi-rational branched covering map. Then F is CLH-equivalent to a sub-hyperbolic rational map G if and only if F has no Thurston obstructions.*

Two different proofs of Theorem 6.1 are provided in [4] and [26]. In the following we will borrow the idea in the proof in [26]. Let us first recall the idea of the proof in [26]. Let T_F denote the Teichmuller space modeled on $(\widehat{\mathbb{C}}, P_F \cup \cup_i \overline{U_i})$. The branch covering F induces an analytic map $\sigma_F : T_F \to T_F$. The existence of G is equivalent to the existence of a fixed point of T_F. The proof in [26] is divided into two steps. In the first step it proves that the non-existence of Thurston obstructions implies certain bounded geometry condition, which will be described later. The second step is to prove the bounded geometry implies the existence of a fixed point.

Now let us describe the bounded geometry condition. Let μ_0 be the standard complex structure. Let μ_n be the complex structure which is the pull back of μ_0 by F^n. Let ϕ_n be the quasiconformal homeomorphism which solves the Beltrami equation given by μ_n and fixes 0, 1 and ∞. Consider the hyperbolic Riemann surface

$$X_n = \widehat{\mathbb{C}} \setminus \phi_n(P_F \cup \cup_i \overline{U_i}).$$

For any non-peripheral curve $\gamma \subset \widehat{\mathbb{C}} \setminus (P_F \cup \cup_i \overline{U_i})$, there is a unique simple closed geodesic η in X_n which is homotopic to $\phi_n(\gamma)$. Let $[\mu_n]$ denote the Teichmüller class of μ_n in T_F and $l_{[\mu_n]}(\gamma)$ the length of η with respect to the hyperbolic metric in X_n. We say γ is a $[\mu_n]$-geodesic if $\phi_n(\gamma)$ is a geodesic in X_n.

Definition 6.2. We say F has bounded geometry if there exists a $\delta > 0$ such that for every non-peripheral curve $\gamma \subset \widehat{\mathbb{C}} \setminus (P_F \cup \cup_i \overline{U_i})$ and all $n \geq 0$, one has $l_{[\mu_n]}(\gamma) > \delta$.

In [3], Chen and Jiang proved that if F does not satisfy the above bounded geometry condition, then there exists at least one non-peripheral curve γ such that $l_{[\mu_n]}(\gamma) \to 0$. Furthermore, they showed that the multi-curve Γ which represents the homotopy classes of all such curves is a Thurston obstruction. Such Thurston obstruction is called the canonical Thurston obstruction.

6.2. **Proof of Theorem 2.1.** Let $f \in \Sigma_{\text{top}}^{\alpha,d}$. Suppose f has no Thurston obstructions in the exterior of Δ. Define

$$F(z) = \begin{cases} f(z) \text{ for } |z| \geq 1 \\ [f(z^*)]^* \text{ for } |z| < 1. \end{cases}$$

Note that $F|\mathbb{T}$ is the rigid rotation $z \mapsto e^{2\pi i\alpha}z$ and $F : \widehat{\mathbb{C}} \to \widehat{\mathbb{C}}$ is a branched covering map of degree $2d-1$ and is symmetric with respect to $\mathbb{T}$. Let $\alpha_n = p_n/q_n$ be a sequence of rational numbers such that $\alpha_n \to \alpha$ as $n \to \infty$. By assumption, $1 \in \Omega_f$. It follows that $1 \in \Omega_F$. Let

$$H = \{z \mid (1+\epsilon)^{-1} < |z| < 1+\epsilon\}$$

be such that

$$(H - \mathbb{T}) \cap (\Omega_F \cup P_F) = \emptyset.$$

For $w \in \widehat{\mathbb{C}}$, let w^* denote the symmetric image of w about $\mathbb{T}$. We can perturb F in H to get a sequence of branched covering maps F_n of the sphere to itself such that

1. $1 \in \Omega_{F_n}$ and $F_n(z^*) = [F_n(z)]^*$,
2. $F_n(z) = F(z)$ for all $z \notin H$,
3. $F_n(z) = e^{2\pi i\alpha_n}z$ for all $z \in \mathbb{T}$,
4. $(H - \mathbb{T}) \cap (\Omega_F \cup P_F) = \emptyset$,
5. all the critical points of F_n in $\mathbb{T}$ belong to the forward orbit of 1, which is denoted as $\mathcal{O}_n$,
6. $F_n \to F$ uniformly with respect to the spherical metric.

Remark 6.1. Since topological branched coverings are quite robust, one shall easily make such perturbations of F. Since $F_n \to F$ uniformly with respect to the spherical metric by (6), the following property holds. For all n large enough, the degree of F_n and the numbers of the critical points of F_n in $\mathbb{T}$, and not in $\mathbb{T}$, are respectively the same as those of F. Moreover, for each critical point c of F, there is exactly one critical point of F_n, say c_n, such that $c_n \to c$ as $n \to \infty$, and the local degree of F_n at c_n is the same as that of F at c.

Lemma 6.1. *F_n has no Thurston obstructions for all n large enough.*

Proof. Our proof is by contradiction. Suppose F_n has a Thurston obstruction for some n large. By [3], F_n would have a canonical Thurston obstruction Γ. Since F_n is symmetric about $\mathbb{T}$, we can make Γ symmetric also, that is, $\gamma \in \Gamma$ if and only if $\gamma^* \in \Gamma$. We claim that all curves in Γ intersect $\mathbb{T}$. Let us prove the claim now. Let $\Gamma' \subset \Gamma$ be the subset consisting of all curves which do not intersect $\mathbb{T}$ and assume $\Gamma' \neq \emptyset$. Since F_n maps $\mathbb{T}$ to $\mathbb{T}$, any non-peripheral component of $F_n^{-1}(\gamma)$ must be disjoint with $\mathbb{T}$ and is homotopic to some element $\eta \in \Gamma$. Note that any curve in $\Gamma \setminus \Gamma'$ is symmetric about $\mathbb{T}$ and can not be homotopic to a non-peripheral curve which does not intersect $\mathbb{T}$. So $\eta \in \Gamma'$. This implies that Γ' is stable. Since $l_n(\gamma) \to 0$ for all $\gamma \in \Gamma'$, Γ' is also a Thurston obstruction. We further divide Γ' into two groups of curves Γ_1 and Γ_2, where Γ_1 consists of all curves in the exterior and Γ_2 the interior. Let us now show that both Γ_1 and Γ_2 are stable. By symmetry it suffices to prove that Γ_2 is stable. Suppose Γ_2 is not stable. Then Γ_2 would contains a curve Γ such that one of the non-peripheral components of $F_n^{-1}(\gamma)$, say η, is homotopic to some ξ in Γ_1. Thus η encloses at least two points of P_{F_n} in the exterior of Δ. By the construction of F_n and F, these points must also belong to P_F and thus P_f. Note that in the exterior of Δ, F_n behaviors like f and maps the inside of η to the inside of γ. In particular, F_n maps these points to the interior of Δ. But This is a contradiction since the action of F_n on these points the same as f and thus maps these points to some points in the exterior of Δ. This contradiction implies that Γ_2 is stable. Thus Γ_1 is stable also. Since $\Gamma' = \Gamma_1 \cup \Gamma_2$ is a Thurston obstruction, by symmetry and the fact that both Γ_1 and Γ_2 are stable, it follows that both Γ_1 and Γ_2 are Thurston obstructions of F_n. Since F_n behaves like f in the exterior of Δ, it follows that Γ_1 is a Thurston obstruction of f. This is a contradiction. This means that $\Gamma' = \emptyset$ and all the curves in Γ intersect $\mathbb{T}$.

Now let μ_0 be the standard complex structure. Let μ_k be the pull back of μ_0 by F_n^k. Let ϕ_k be the qc homeomorphism of the sphere which fixes 0, 1 and ∞ and solves the Bletrami equation given by μ_k. Then

$$G_n^k = \phi_{k-1} \circ F_n \circ \phi_k^{-1}, \;\; k \ge 1$$

are rational maps and thus Blaschke products by symmetry.

First we note that the sequence $\{G_n^k = \phi_k^{-1} \circ F_{n,m} \circ \phi_{k-1}\}$ is a sequence of Blaschke products of $2d-1$ whose restrictions to $\mathbb{T}$ are circle homeomorphisms. By a lemma of Herman (cf. §15 of [9]), there is a uniform $\kappa(d) > 0$ depending only on d such that $\{G_n^k\}$ is a normal family of holomorphic functions in a κ_0-neighborhood of $\mathbb{T}$. This implies

Fact 1. The sequence $\{G_n^k\}$ is equicontinuous in a κ_0-neighborhood of $\mathbb{T}$.

We claim that every γ in Γ encloses at least a pair of points in $P_{F_n} \setminus \mathcal{O}_n$, which is symmetric about $\mathbb{T}$. If not, by symmetry, γ enclose no point in $P_{F_n} \setminus \mathcal{O}_n$. Since γ is non-peripheral, it would enclose two points in $\mathcal{O}_n$. Since $l_{[\mu_k]}(\gamma) \to 0$ as $k \to \infty$, to get a contradiction, it suffices to show that there is a $d_0 > 0$ depending only on n such that for any two distinct points $x, y \in \mathcal{O}_n$,

$$d(\phi_k(x), \phi_k(y)) > d_0, \;\; \forall k \ge 1, \tag{45}$$

where $d(x, y)$ denotes the Euclidean length of the smaller arc connecting x and y. We prove (45) by contradiction. First note that $d_T([\phi_k], [\phi_{k-1}]) \le d_T([\phi_1], [\phi_0])$ for all $k \ge 1$ (The inequality comes from the fact that the Thurston pull back does not increase the Teicmuller distance, see [8]). This, together with Proposition 7.2 of [8], implies that

$$e^{-2} \cdot l_{[\mu_{k-1}]}(\gamma) \le l_{[\mu_k]}(\gamma) \le e^2 \cdot l_{[\mu_{k-1}]}(\gamma). \tag{46}$$

Because all ϕ_k fix 0, 1 and ∞, the above inequality implies the following fact.

Fact 2. There is an $\epsilon > 0$ such that for any two integers $1 \le k' \le k_2 < k' + q_n$, and any two points a and b in $\mathcal{O}_n$, if $d(\phi_{k'}(a), \phi_{k'}(b)) < \epsilon$, then $d(\phi_{k''}(a), \phi_{k''}(b)) < 2\pi/q_n$.

Now assume x and y are two adjacent points in $\mathcal{O}_n$ and $d(\phi_N(x), \phi_N(y))$ is small. Since $\mathcal{O}_n$ is a periodic orbit of F_n of period q_n, there is some $1 \le l \le q_n$ such that $d(\phi_N(F_n^l(x)), \phi_N(F_n^l(y))) \ge 2\pi/q_n$. But on the other hand, since $\{G_n^k\}$ is a normal family of holomorphic functions in the $\kappa(d)$-neighborhood of $\mathbb{T}$, they are equicontinuous in $\mathbb{T}$. We thus have a $d_0 > 0$ depending only on n such that

$$d(\phi_{N-l}(F_n^l(x)), \phi_{N-l}(F_n^l(y))$$

$$= d(G_{N-l} \circ G_{N-l+1} \circ \cdots G_{N-1}(\phi_N(x)), G_{N-l} \circ G_{N-l+1} \circ \cdots G_{N-1}(\phi_N(y))) < \epsilon$$

provided that $d(\phi_N(x), \phi_N(y)) \le d_0$. For such $d_0 > 0$, if $d(\phi_N(x), \phi_N(y)) \le d_0$, we would have $d(\phi_{N-l}(F_n^l(x), \phi_{N-l}(F_n^l(y)) < \epsilon$. By the Fact 2 above we then have $d(\phi_N(F_n^l(x)), \phi_N(F_n^l(y))) < 2\pi/q_n$. This is a contradiction. This proves the claim.

From the claim we just proved, it follows that γ encloses (at least) a pair of points in $P_{F_n} \setminus \mathcal{O}_n$, which are symmetric about $\mathbb{T}$. Let us denote this pair of points as z and z^*. Since $l_{[\mu_k]}(\gamma) \to 0$, it follows that $d(\phi_k(z), \mathbb{T}) = d(\phi_k(z^*), \mathbb{T}) \to 0$ as $k \to \infty$. Note that the forward orbits of z and z^* eventually land on two holomorphic attracting periodic cycles of F_n. Suppose $w = F_n^l(z)$ and $w^* = F_n^l(z^*)$ belongs to a holomorphic attracting cycle of F_n. By Fact 1, it follows that $d(\phi_k(w), \mathbb{T}) = d(\phi_k(w^*), \mathbb{T}) \to 0$ as $k \to \infty$. This implies there is a $\eta \in \Gamma$ which encloses w and w^*. Assume that the period of w is p. Let $I_j, j = 1, \cdots, q_n$, denote the components of $\mathbb{T} \setminus \mathcal{O}_n$. Take N large. Let I_i is the one such that

$$d(\phi_N(I_i), \phi_N(w)) = \min_{1 \le l \le q_n} \{d(\phi_N(I_l), \phi_N(w))\}.$$

Let η' be a $[\mu]_N$-geodesic in the homotopic class of η. Then η' intersects either I_i or one of the adjacent components of I_i. Let $I_j = F_n^p(I_i)$. From Fact 1 it follows that $\phi_{N-p}(I_j)$ is close to $w = F_n^p(w)$ also. This implies there exists a $[\mu]_{N-p}$-geodesic ζ' which is homotopic to some element in Γ, say ζ, which intersects I_j or one of the adjacent intervals of I_j and encloses w and w^* in its inside. Let S and R denote the two arcs connecting I_i and I_j. Since α is irrational, when n is large enough, both R and S contains at least two components of $\mathbb{T} \setminus \mathcal{O}_n$. This implies that any curve in the homotopic class of η' must intersect a curve in the homotopy class of ζ. In particular, η intersects ζ. This is a contradiction. The proof of Lemma 6.1 is completed. □

By Lemma 6.1 and Theorem 6.1, it follows that F_n is CLH-equivalent a rational map G_n. By symmetry, G_n can be taken to be a Blaschke product.

Lemma 6.2. *The sequence $\{G_n\}$ is pre-compact.*

Proof. Note that all the finite poles of G_n belong to the interior of Δ and except the origin, all the zeros of G_n belong to the exterior of Δ. By a lemma of Herman (cf. §15 of [9]), these zeros and poles are uniformly bounded away from $\mathbb{T}$. So to prove the lemma, it suffices to show that all the zeros of G_n are uniformly bounded. Note that

$$G_n(z) = \lambda_n z^d \prod_{i=1}^{d-1} \frac{z - p_i}{1 - \overline{p_i} z} \tag{47}$$

where $|\lambda_n| = 1$ and $|p_i| > 1$, $1 \le i \le d-1$. Let

$$\epsilon = (-1)^{d-1} \prod_{i=1}^{d-1} \frac{1}{\overline{p_i}}.$$

Then near infinity $G_n(z) = \epsilon z^d + o(z^d)$. Take η such that $\eta^{d-1} = \epsilon$. Consider the map

$$\tilde{G}_n(\zeta) = \eta \cdot G_n(\frac{\zeta}{\eta}).$$

Then $\tilde{G}_n(\zeta) = \zeta^d + o(\zeta^d)$ near infinity. Let Φ be a holomorphic map in a neighborhood of ∞ such that $\Phi(z) = z(1 + o(1))$ near infinity which conjugates $\tilde{G}_n$ to $z \mapsto \zeta^d$. Since all critical orbits of G_n, and thus of $\tilde{G}_n$, are bounded, it follows that Φ can be extended to a Riemann isomorphism between the exterior of the unit disk to the immediate basin of ∞ for $\tilde{G}_n$. By Koebe's 1/4-theorem, the immediate attracting basin of $\tilde{G}_n$ at infinity contains the exterior of $\{z \mid |z| < 4\}$. Go back to G_n, we get

$$|p_i| \le \frac{4}{|\eta|}, \ \ 1 \le i \le d-1.$$

This implies

$$\min_{1 \le i \le d-1} |p_i| \le \max_{1 \le i \le d-1} |p_i| \le 4 \min_{1 \le i \le d-1} |p_i|.$$

From the above inequality, it follows that $|p_i|$ are all large provided that one of them is large. From (47), it then follows that G_n is holomorphic in $1/2 < |z| < 2$, and moreover, G_n can be arbitrarily close to $\lambda_n z$ in H provided that these $|p_i|$ are large enough. But this would imply G_n has no critical point in $\mathbb{T}$. This is a contradiction. The proof of Lemma 6.2 is completed.

□

Remark 6.2. From the proof of Lemma 6.2, there is a constant $R(d) > 1$ such that $|p_i| < R(d)$ for all G_n. From this it follows that the immediate attracting basin at infinity for G_n contains the exterior of $\{z \, |z| \le 4R(d)\}$.

By taking a subsequence, we may assume that G_n converges to G uniformly with respect to the spherical metric. Then $G|\mathbb{T}$ is a critical circle homeomorphism with rotation number α. So there is a qs circle homeomorphism h which conjugates $G|\mathbb{T}$ to R_α. Let ϕ_n and ψ_n be the pair of homeomorphisms of the sphere such that

$$\phi_n \circ F_n = G_n \circ \psi_n$$

and ϕ_n is isotopic to ϕ_n rel P_{F_n}, moreover, $\phi_n|D_i = \psi_n|D_i$ are holomorphic.

Proposition 6.1. There exist a pair of homeomorphisms of the sphere ϕ and ψ which fix 0, 1 and ∞ such that

1. $F \circ \phi = G \circ \psi$,
2. ϕ is isotopic to ψ rel $\mathbb{T} \cup P \cup \cup_i D_i$,
3. $\phi = \psi$ is holomorphic on each D_i.

The proof of Proposition 6.1 is divided into several lemmas.

Lemma 6.3. *$\phi_n|\mathbb{T} \to h$ and $\psi_n|\mathbb{T} \to h$ uniformly.*

Proof. All the maps in the proof are considered only on $\mathbb{T}$. So to simplify the notations, we just write $\phi_n|\mathbb{T}$ and $\psi_n|\mathbb{T}$ as ϕ_n and ψ_n respectively. It suffices to prove ϕ_n and ψ_n converge uniformly. Let ϕ and ψ be the limit maps respectively. Since the convergence is uniform, it follows that both ϕ and ψ are circle homeomorphisms. For any $k \ge 0$ we have $\phi_n(F_n^k(1)) = \psi_n(F_n^k(1)) = G_n^k(1)$. Since $F_n \to F$ and $G_n \to G$ uniformly on $\mathbb{T}$, let $n \to \infty$ we get $\phi(F^k(1)) = \psi(F^k(1)) = G^k(1)$ for all $k \ge 1$. This implies that ϕ and ψ coincide on $\{F^k(1)\}_{k=0}^{\infty}$, which is a dense subset of $\mathbb{T}$. It follows that $\phi = \psi$. Since $\phi(1) = \psi(1) = h(1) = 1$, it follows that $\phi = \psi = h$. In the following let us only prove that ϕ_n converges uniformly. The same method will work for ψ_n.

By construction, the point 1 is a critical point for F, G, F_n and G_n. For any $N \ge 1$, since $F_n \to F$ and $G_n \to G$ uniformly, by taking n large enough, we can make sure the orbit segments

$$\mathcal{O}_N(G) = \{1, G^1(1), \cdots, G^N(1)\} \text{ and } \mathcal{O}_N(G_n) = \{1, G_n^1(1), \cdots, G_n^N(1)\}$$

have the same order, and

$$\mathcal{O}_N(F) = \{1, F^1(1), \cdots, F^N(1)\} \text{ and } \mathcal{O}_N(F_n) = \{1, F_n^1(1), \cdots, F_n^N(1)\}$$

have the same order. Since ϕ_n and ψ_n are circle homeomorphisms and map $\mathcal{O}_N(F_n)$ to $\mathcal{O}_N(G_n)$ with the order being preserved, thus $\mathcal{O}_N(F_n)$ and $\mathcal{O}_N(G_n)$ have the same order. All of these implies that for any fixed N, by taking n large enough, all the four orbit segments, $\mathcal{O}_N(F)$, $\mathcal{O}_N(F_n)$, $\mathcal{O}_N(G)$ and $\mathcal{O}_N(G_n)$, have the same order.

Let $\epsilon > 0$ be an arbitrary small number. Since $G|\mathbb{T}$ and $F|\mathbb{T}$ are circle homeomorphisms with irrational rotation number, by taking N large enough we can make sure the length of each component of $\mathbb{T} \setminus \mathcal{O}_N(F)$ and $\mathbb{T} \setminus \mathcal{O}_N(G)$ is less than $\epsilon/4$. Since $F_n \to F$ and $G_n \to G$ uniformly, there exists an $M' > 1$ such that for all $n \ge M'$, the length of each component of $\mathbb{T} \setminus \mathcal{O}_N(F_n)$ and $\mathbb{T} \setminus \mathcal{O}_N(G_n)$ is less than $\epsilon/3$. For a component I of $\mathbb{T}\setminus\mathcal{O}_N(F_n)$, we use I^l and I^r to denote the components of $\mathbb{T}\setminus\mathcal{O}_N(F_n)$, which are adjacent to I from the left and right respectively. Similarly, for a component J of $\mathbb{T} \setminus \mathcal{O}_N(G_n)$, we use J^l and J^r to denote the components of $\mathbb{T} \setminus \mathcal{O}_N(G_n)$, which are adjacent to J from the left and right respectively. For N given, since $F_n \to F$ and $G_n \to G$ uniformly on $\mathbb{T}$, there exists an $M > M'$ such that for all $m, n \ge M$, the following holds: For any component I of $\mathbb{T} \setminus \mathcal{O}_N(F_n)$ and any component J of $\mathbb{T} \setminus \mathcal{O}_N(G_n)$, let $\tilde{I}$ and $\tilde{J}$ be the corresponding component of $\mathbb{T} \setminus \mathcal{O}_N(F_m)$ and $\mathbb{T} \setminus \mathcal{O}_N(G_m)$ respectively, then $I \subset \overline{\tilde{I}^l \cup \tilde{I} \cup \tilde{I}^r}$ and $J \subset \overline{\tilde{J}^l \cup \tilde{J} \cup \tilde{J}^r}$.

Now let $z \in \mathbb{T}$ be an arbitrary point and I be a component of $\mathbb{T} \setminus \mathcal{O}_N(F_n)$ such that $z \in \overline{I}$. We then have $z \in \overline{\tilde{I}^l \cup \tilde{I} \cup \tilde{I}^r}$. For the above M, let $n, m > M$ be any two integers. Let $J = \phi_n(I)$. Then $\tilde{J} = \phi_m(\tilde{I})$. It follows that

$$\phi_n(z) \in \overline{J} \subset \overline{\tilde{J}^l \cup \tilde{J} \cup \tilde{J}^r}.$$

and

$$\phi_m(z) \in \phi_m(\overline{\tilde{I}^l \cup \tilde{I} \cup \tilde{I}^r}) = \overline{\tilde{J}^l \cup \tilde{J} \cup \tilde{J}^r}.$$

Then $|\phi_n(z) - \phi_m(z)| \le |\tilde{J}^l| + |\tilde{J}| + |\tilde{J}^r| < \epsilon$. This proves that ϕ_n converges uniformly. Lemma 6.3 then follows. ☐

Let

$$P = P_{F_n} \setminus (\mathbb{T} \cup \cup_i D_i) = P_F \setminus (\mathbb{T} \cup \cup_i D_i).$$

Then P is a finite set.

Lemma 6.4. *There exist a $K > 1$ such that for all $n \ge 1$,*

1. $\phi_n(P \cup \cup_i D_i) \subset \{z \mid 1/K < |z| < K\}$,
2. *for every holomorphic disk D_i,* $\operatorname{diam}(\phi_n(D_i)) > 1/K$.
3. *for every two distinct points z and w in P,* $\operatorname{dist}(\phi_n(z), \phi_n(w)) > 1/K$,
4. *for every point $z \in P$ and every holomorphic disk D_i,* $\operatorname{dist}(\phi_n(z), \phi_n(D_i)) > 1/K$,
5. *for every holomorphic disk D_i,* $\operatorname{dist}(\phi_n(D_i), \mathbb{T}) > 1/K$,
6. *for every two distinct holomorphic disks D_i and D_j,* $\operatorname{dist}(\phi_n(D_i), \phi_n(D_j)) > 1/K$.
7. *for every point $z \in P$,* $\operatorname{dist}(\phi_n(z), \mathbb{T}) > 1/K$,

Proof. By Remark 6.2 it follows that there is a $K > 1$ such that $\phi_n(P \cup \cup_i D_i) \subset \{z \,|\, |z| < K\}$. By symmetry we further have $\phi_n(P \cup \cup_i D_i) \subset \{z \mid 1/K < |z| < K\}$. This is the property (1).

Now let us prove (2). Let us choose a D_i. For all n large, let ϕ_n and ψ_n be the homeomorphisms of the sphere in Definition 6.1 such that $\phi_n \circ F_n = G_n \circ \psi_n$. Then $\phi_n(D_i)$ is contained in the immediate attracting basin of some attracting cycle of G_n. Let Ω_n denote this basin. Note that all G_n are combinatorially equivalent when restricted to this immediate attracting basin. Since such equivalence is locally holomorphic, it is actually a holomorphic conjugation. This implies that $\phi_n(D_i)$ can not be small. This proves (2).

To show (3), it suffices to show the case that both z and w belong to holomorphic attracting disks. Because otherwise we can consider the forward images of z and w by F_n^k for some k large enough. But then it is obvious.

The property (4), (5), (6) follow from the fact that D_i has a collar neighborhood on which ϕ_n is holomorphic.

To prove (7) note that G_n preserve $\mathbb{T}$ and the sequence $\{G_n\}$ is equicontinuous. But on the other hand, $G_n^k(z)$ belongs to some $\phi_n(D_i)$ for some $k > 0$ which is bounded away from $\mathbb{T}$. This implies (7).

The proof of Lemma 6.4 is completed. □

By (1) and (2) of Lemma 6.4, the sequence $\{\phi_n | D_i = \psi_n | D_i\}$ is precompact. By taking a subsequence if necessary we may assume that $\phi_n | D_i$ converge uniformly to some univalent map on D_i. Let us denote the limit of $\phi_n(U_i)$ by U_i. Let

$$Q = P_G \setminus (\mathbb{T} \setminus \cup_i U_i).$$

From Lemma 6.4 it follows that

Lemma 6.5. *When restricted to* $P \cup \cup_i D_i$, $\phi_n = \psi_n$ *converge uniformly to some injective map which maps* P *to* Q *and maps each* D_i *to* U_i.

Let us now define a map σ on $\mathbb{T} \cup P \cup \cup_i D_i$. On $\mathbb{T}$, define $\sigma = h$. On $P \cup \cup_i D_i$, define σ to be the limit map in Lemma 6.5. Note that σ maps critical values of F to the critical values of G.

Let us now begin the proof of Proposition 6.1.

Choose n large enough such that F_n is close to F and G_n is close to G. Let $\eta : \widehat{\mathbb{C}} \to \widehat{\mathbb{C}}$ be a homeomorphism of the sphere which is nearby the identity map such that

1. η is the identity map in the exterior of a small annular neighborhood of the unit circle,
2. η is symmetric about the unit circle, and in particular, it preserve the unit circle,
3. η maps each critical value of F to the corresponding critical value of F_n.

Note that $P_F \setminus \mathbb{T} = P_{F_n} \setminus \mathbb{T}$, so to get η, one need only perturb the identity map symmetrically in a small neighborhood of the unit circle so that it maps each critical value of F in $\mathbb{T}$ to the corresponding one of F_n. It is clear that η can be lifted to a homeomorphism $\tilde{\eta}$ such that the following diagram commutes. Moreover, $\tilde{\eta}$ can be arbitrarily close to the identity map provided that η is close to the identity map enough.

$$\begin{tikzcd} \widehat{\mathbb{C}} \arrow[r, "\tilde{\eta}"] \arrow[d, "F"'] & \widehat{\mathbb{C}} \arrow[d, "F_n"] \\ \widehat{\mathbb{C}} \arrow[r, "\eta"] & \widehat{\mathbb{C}} \end{tikzcd}$$

Similarly, let $\xi : \widehat{\mathbb{C}} \to \widehat{\mathbb{C}}$ be a homeomorphism of the sphere which is nearby the identity map and is symmetric about the unit circle, and moreover,

$$\xi = \sigma \circ \eta^{-1} \circ \phi_n^{-1} \text{ on } \mathbb{T} \cup \phi_n(P \cup \cup_i D_i).$$

It is clear that ξ maps each critical value of G_n to the corresponding critical value of G. Then ξ can be lifted to a homeomorphism $\tilde{\xi}$ such that the following diagram commutes. As before, $\tilde{\xi}$ can be arbitrarily close to the identity map provided that ξ is close to the identity map enough.

$$\begin{tikzcd} \widehat{\mathbb{C}} \arrow[r, "\tilde{\xi}"] \arrow[d, "G_n"'] & \widehat{\mathbb{C}} \arrow[d, "G"] \\ \widehat{\mathbb{C}} \arrow[r, "\xi"] & \widehat{\mathbb{C}} \end{tikzcd}$$

By Lemma 6.3 we may assume that $\phi_n|\mathbb{T}$ is close to h by taking n large enough. Now define

$$\phi = \xi \circ \phi_n \circ \eta \text{ and } \psi = \tilde{\xi} \circ \psi_n \circ \tilde{\eta}.$$

By the construction it follows that $\phi = \psi = \sigma$ on $\mathbb{T} \cup P \cup \cup D_i$. That is,

$$\phi|\mathbb{T} = \psi|\mathbb{T} = h,$$

and ϕ maps each point in P to the corresponding point in Q, and maps each D_i to U_i.

From the above construction it follows that all the maps $\eta, \tilde{\eta}, \xi$, and $\tilde{\xi}$ can be arbitrarily close to the identity map provided that n is large enough. Thus ϕ_n and ψ_n can be arbitrarily close to ϕ and ψ provided that n is large enough. Since both $\{\mathbb{T}, P, \text{ all the } D_i\}$ and $\{\mathbb{T}, Q, \text{ all the } U_i\}$ contain finitely many objects and $\phi = \psi$ maps each object in the first group homeomorphically to the corresponding one in the second group, the isotopy class of ϕ and ψ rel $\mathbb{T} \cup P \cup \cup_i D_i$ are determined respectively by ϕ_n and ψ_n provided that n is large enough. Since ϕ_n is isotopic to ψ_n rel $\mathbb{T} \cup P \cup \cup D_i$ for all n, we can make sure ϕ and ψ constructed in this way are isotopic to each other rel $\mathbb{T} \cup P \cup \cup_i U_i$ by taking a large n in the beginning of the construction. This completes the proof of Proposition 6.1.

Now we can prove the existence part of Theorem 2.1 by performing qc surgery on G. The process is routine. Let $H : \Delta \to \Delta$ be a qc extension of h which fixes the origin. Define

$$\widehat{G}(z) = \begin{cases} G(z) \text{ for } |z| \geq 1 \\ H^{-1} \circ R_\theta \circ H(z) \text{ for } |z| < 1 \end{cases} \tag{48}$$

Let μ_0 be the pull back of the standard complex structure by H. Then pull back μ_0 to the whole plane by the iteration of $\widehat{G}$ and get a $\widehat{G}$-invariant complex structure μ. Let ϕ be the qc homeomorphism of the plane to itself which solves the Beltrmai equation given by μ and fixes 0 and 1. It is clear that $g = \phi \circ \widehat{G} \circ \phi^{-1}$ is a Siegel polynomial map in $\Sigma_{\text{geom}}^{\alpha,d}$. Note that in the immediate basin of each attracting cycle of G which lies in the exterior of Δ, $\mu = 0$ and thus ϕ is holomorphic. This, together with the fact that $F = f$ in the exterior of Δ, and that G is CLH-equivalent in the sense of Lemma 6.1, implies that g is CLH-equivalent to f. This proves the existence part of Theorem 2.1.

Now it remains to prove the uniqueness part. Suppose $f \in \Sigma_{\text{top}}^{\alpha,d}$ is CLH-equivalent to two maps σ and ξ in $\Sigma_{\text{geom}}^{\alpha,d}$. We will show that σ is linearly conjugate to ξ. To see this, let G, $\widehat{G}$ and g be maps obtained in the proof of the existence part. Let $K_{\widehat{G}}$ be the set of all points whose forward orbits under the iteration of $\widehat{G}$ is bounded. Let $J_{\widehat{G}} = \partial K_{\widehat{G}}$. We call $J_{\widehat{G}}$ the Julia set of $\widehat{G}$.

Lemma 6.6. *The set* $J_{\widehat{G}}$ *has zero Lebesgue measure.*

Before we prove the lemma, let us show how it implies the uniqueness part of Theorem 2.1. Indeed, if Lemma 6.6 holds, then J_g has zero Lebesgue measure. Since f is CLH-equivalent to g and σ, and since the boundary of the Siegel disks of both g and σ are quasi-circles, g and σ are qc-equivalent to each other, and moreover, the qc-equivalence can be made to be a holomorphic conjugation when restricted to the Siegel disk and an open neighborhood of all the attracting cycles. That is, we have a pair of qc homeomorphisms ϕ_1 and ϕ_2 such that

1. $\phi_1 \circ g = \sigma \circ \phi_2$,
2. ϕ_1 is isotopic to ϕ_2 rel P_g,
3. $\phi_1 = \phi_2$ is holomorphic in the union of the interior of the Siegel disk and an open neighborhood of all attracting periodic cycles of g.

By induction we get a sequence of qc homeomorphisms ϕ_{k+1} by lifting ϕ_k through

$$\phi_k \circ g = \sigma \circ \phi_{k+1}.$$

Note that the qc constants of ϕ_k is bounded that of ϕ_1. Let μ_k be the Beltrami coefficient of ϕ_k. Since all the points in the Fatou set is either contracted to some attracting cycle of g, or is eventually mapped to the Siegel disk of g, so $\mu_k \to 0$ on the Fatou set of g. Since J_g is the qc image of $J_{\widehat{G}}$, by Lemma 6.6 J_g has zero Lebesgure measure. So $\mu_k \to 0$ a.e. This implies that ϕ_k converges to some linear map. In particular, g is linearly conjugate to σ. Using the same way we can show that g is linearly conjugate to ξ. This implies that σ is linearly conjugate to ξ. This proves the uniqueness part of Theorem 2.1.

Now it remains to prove Lemma 6.6. The proof is by contradiction. Suppose $J_{\widehat{G}}$ has positive Lebesgue measure. Let z_0 be a Lebesgue point of $J_{\widehat{G}}$. For $n \geq 1$ let $z_n = G^n(z_0)$. By a standard pull back argument (see Proposition of [12]), z_n accumulates to $\mathbb{T}$. The contradiction is derived by using an idea of McMullen [13]. The idea can be sketched as follows. We first show that there is a subsequence n_j for which z_{n_j} pass through the cones spanned at the critical points in $\mathbb{T}$. Then we can take a small disk B_j in the cone such that

1. $G(B_j) \subset \Delta$,
2. $\text{dist}(B_j, \mathbb{T}) \asymp \text{dist}(B_j, z_{n_j}) \asymp \text{dist}(z_{n_j}, \mathbb{T})$.

From (2) we can take a Jordan domain A_j which is disjoint with P_G and contains both B_j and z_{n_j} such that

$$\text{diam}(A_j) \asymp \text{dist}(A_j, \mathbb{T}).$$

Let X be the unbounded component of $\widehat{\mathbb{C}} \setminus P_G$. The above relation implies that

$$\mathrm{diam}_X(A_j) < K$$

for some uniform constant $K > 0$. Now we pull back A_j along the orbit $z_0, \cdots, z_{n_j}$, and denote the component of $G^{-k}(A_j)$ which contains z_{n_j-k} by $A_j^{n_j-k}$. Let X_k be the unbounded component of $G^{-k}(X)$. Then $G^k : X_k \to X$ is a holomorphic covering map and $X_k \subset X$. Note that $A_j^{n_j-k} \subset X_k$. It follows that

$$\mathrm{diam}_X(A_j^{n_j-k}) \le \mathrm{diam}_{X_k}(A_j^{n_j-k}) = \mathrm{diam}_X(A_j) < K.$$

For $1 \le i \le j$, Since $A_j^{n_i}$ contains z_{n_i} and z_{n_i} is contained in some cone spanned at some critical point, it follows that $A_j^{n_i}$ is contained in a definite cone spanned at this critical point. Thus the inverse branch of G^{-1} which maps $A_j^{n_i+1}$ to $A_j^{n_i}$ contracts the hyperbolic diameter by a definite factor $0 < \delta < 1$, that is,

$$\mathrm{diam}_X(A_j^{n_i} < \delta \cdot \mathrm{diam}_X(A_j^{n_i+1}).$$

This implies that $\mathrm{diam}_X(A_j^0) \to 0$ as $j \to \infty$. In addition, since $\mathrm{dist}(A_j, P_G) \asymp \mathrm{diam}(A_j)$, the distortion of G^{-n_j} is uniformly bounded. Let B_0 denote the component of $G^{-n_j}(B)$ which is contained in A_j^0. We have

$$area(B_0) \succeq \mathrm{diam}(A_j)^2.$$

Since B and thus B_0 is disjoint from $J_{\widehat{G}}$, we get a contradiction with the assumption that z_0 is a Lebesgue point of $J_{\widehat{G}}$.

Now we need to show the existence of such sequence n_j. We use a similar argument with the one used in [24]. For each open arc $I \subset \mathbb{T}$, consider the space

$$\Omega_I = \widehat{\mathbb{C}} \setminus (P_G \setminus I).$$

Let $d_{\Omega_I}(\cdot, \cdot)$ denote the hyperbolic distance in Ω_I and

$$H_d(I) = \{z \in \Omega \mid d_{\Omega_I}(z, I) < d\}.$$

As in Lemma 4.18, when I is small, $H_d(I)$ is like the hyperbolic neighborhood in the slit plane, that is, it is almost like the domain bounded two arcs of Euclidean circles which are symmetric about each other and such that the four exterior angles formed by the two arcs and $\mathbb{T}$ are all equal to α with $d = \ln \cot(\alpha/4)$.

Fix a $d > 0$. Let $h : \mathbb{T} \to \mathbb{T}$ be the circle homeomorphism such that $G|\mathbb{T} = h^{-1} \circ R_\alpha \circ h$. For each z_n, let $I_n \subset I$ be the arc such that $z_n \in H_d(I_n)$ and moreover, I_n is the smallest one in the following sense

$$|h(I_n)| = \min_{I \subset I} |h(I)|$$

where the min is taken over all arc $I \subset$ such that $z_n \in H_d(I)$. Since $z_n \to \mathbb{T}$, we have $|I_n| \to 0$ and thus

$$|h(I_n)| \to 0$$

as $n \to \infty$. So there is an increasing subsequence of integers, say m_j, such that

$$|h(m_j)| < |h(I_n)| \quad \text{for all } 1 \le n < m_j.$$

Let $n_j = m_j - 1$. We claim $\{n_j\}$ is the desired subsequence. Let us prove the claim. These are two cases. In the first case, I_{m_j} contains no critical value. In the second case, I_{m_j} contains a critical value. In the first case, let $J \subset \mathbb{T}$ be the arc such that

$G(J) = I_{m_j}$. Let U be the component of $G^{-1}(H_d(I_{m_j}))$ which is attached to J. By Schwarz lemma it follows that $U \subset H_d(J)$. By the minimal property of I_{m_j}, it follows that $z_{m_j-1} = z_{n_j} \notin H_d(J)$. This implies that z_{m_j} is near a critical value in $\mathbb{T}$, and $H_d(J)$ is near some critical point c. By taking $d > 0$ small $U \subset H_d(J)$ is contained in some angle domain bounded by a ray from c and $\mathbb{T}$ with small angle. This implies that z_{n_j} is contained in a definite cone spanned at c. In the second case, I_{n_i} contains a critical point. If the angle between $[c, z_{n_j}]$ is small, we would have a J such that $z_{n_j} \subset H_d(J)$ with $|J| \ll |[c, z_{n_j}] \preceq |I_{n_j}|$ and thus $h(J) < h(I_{n_j})$. This is a contradiction with the minimal property of I_{n_j}. This proves Lemma 6.6. The proof of Theorem 2.1 is completed.

6.3. **Proof of the Key-Lemma 2.** Let $g \in \Sigma^{\alpha,d}_{\text{geom}}$. Let $f \in \Sigma^{\alpha,d}_{\text{top}}$ such that f is CLH-equivalent to g. Then f has no Thurston obstructions by a Theorem of McMullen. By perturbing f in a small neighborhood of the unit disk, we get a sequence $f_n \in \Sigma^{\alpha,d}_{\text{top}}$ such that all the critical points in $\mathbb{T}$ lies in a periodic orbit, and moreover, f_n has no Thurston obstructions. By Theorem 2.1, we get a sequence $g_n \in \Sigma^{\alpha,d}_{\text{geom}}$ which is CLH-equivalent to f_n for all $n \geq 1$.

Recall in the construction of g_n, we first do a symmetrization of f_n and get a branched covering map of the sphere, say F_n. Then we get a Blaschke product G_n which is CLH-equivalent to F_n. That is, there is a pair of homeomorphisms ϕ_n and ψ_n such that $\phi_n \circ F_n = G_n \circ \psi_n$. Lemma 6.4 applies to $\{G_n\}$. Thus $\{G_n\}$ is pre-compact. By taking a subsequence of G_n we may assume that $G_n \to G$. Let F be the symmetrization of f. As in the proof of Proposition 6.1, it follows that F is CLH-equivalent to G. Thus $\phi^{-1} \circ \widehat{G} \circ \phi = g$.

By taking a subsequence we may assume that $\phi_n \to \phi$. This proves the Key-Lemma 2.

7. Appendix

Lemma 7.1. *Let f be a degree-d polynomial map with a Siegel disk centered at the origin. Then there exist $M > 1$ depending only on d such that if f has two critical points c and c' with $|c|/|c'| > M$, then there exist a pair of domains $U \Subset V$ containing the Siegel disk of f centered at the origin such that the tuple (U, V, f) is a polynomial-like map which is qc conjugate to a polynomial map g of degree less than d. In particular, there exists a $L > 1$ which depends only on d such that*

$$\operatorname{diam}(D) \leq L \cdot \min_{c \in \Omega_f} |c|$$

where D is the Siegel disk of f centered at the origin and Ω_f is the set of all critical points of f.

Proof. Let $c_1, \cdots, c_{d-1}$ be the critical points of f. Through a linear conjugation, we may assume that $1 = |c_{d-1}| \leq |c_{d-2}| \leq \cdots \leq |c_1|$.

Claim. For each $M_l > 1$, there exists an $M_{l+1} > M_l$ such that if

$$1 = |c_{d-1}| \leq |c_{d-2}| \leq \cdots \leq |c_{d-l}| \leq M_l < M_{l+1} < |c_{d-l-1}| \leq \cdots \leq |c_1|, \tag{49}$$

then there exist domains $U \Subset V$ containing the origin such that (f, U, V) is a polynomial map which is qc conjugate to some polynomial of degree $l+1$. Let us prove the Claim first. Recall that

$$f(z) = f_{c_1,\cdots,c_{d-1}}(z) = \sum_{i=1}^{d} a_i z^i$$

with

$$a_i = e^{2\pi i\alpha} \cdot \left(\frac{(-1)^{i-1}}{i}\right) \cdot \frac{Q_{d-i}(c_1,\cdots,c_{d-1})}{c_1\cdots c_{d-1}}$$

where Q_{d-i} is the degree-$(d-i)$ elementary polynomials of $c_1,\cdots,c_{d-1}$. If (49) holds, a simple calculation shows that there is some constant

$$C = C(d, M_l) > 1$$

depending only on d and M_l such that by taking M_{l+1} large enough, the following inequalities hold.

(i) $C^{-1} < |a_{l+1}| < C$,
(ii) $|a_k| \le C$ for $1 \le k \le l$,
(iii) $|a_k| < d! \cdot M_{l+1}^{-1}$ for $l+2 \le k \le d$.

From (i) and (ii) it follows that by taking $R > 0$ large enough, we can make sure that

$$\sum_{i=1}^{l} |a_i||z|^i \ll |a_{l+1}z^{l+1}| \quad \text{for } |z| \ge \left(\frac{R}{2C}\right)^{1/(l+1)}. \tag{50}$$

Fix such an R. Then from (iii) it follows that if we take

$$M_{l+1} > R$$

large enough, then we have

$$\sum_{i=l+2}^{d} |a_i||z|^i \ll |a_{l+1}z^{l+1}| \quad \text{for } |z| \le |C\cdot R^{l+1}|. \tag{51}$$

Now define a quasi-regular map F as follows.

$$F(z) = \begin{cases} f(z) \text{ for } |z| \le R \\ f(z) - \frac{|z|-R}{|a_{l+1}R^{l+1}|-R}(f(z) - a_{l+1}z^{l+1}), \text{ for } R < |z| < |a_{l+1}R^{l+1}| \\ a_{l+1}z^{l+1} \text{ if } |z| \ge |a_{l+1}R^{l+1}|. \end{cases}$$

By definition, F is holomorphic for $|z| < R$ and $|z| > |a_{l+1}R^{l+1}|$. From (i-iii) and a simple calculation we get

$$|F_{\bar z}| \ll |F_z| \text{ for } R < |z| < |a_{l+1}R^{l+1}|.$$

provided that R and M_{l+1} are large enough (the choice of M_{l+1} depends on R). This implies that the real dilatation of F in $\{z\,|R < |z| < |a_{l+1}R^{l+1}|\}$ can be arbitrarily close to 1. Note that the forward orbit of any point z passes through the annulus $\{z\,|R \le |z| \le |a_{l+1}R^{l+1}|\}$ at most two times. By a routine argument it follows that F is conjugate to a polynomial of degree $l+1$ through some K-qc homeomorphism where $K > 1$ can be arbitrarily close to 1 provided that R and M_{l+1} are chosen appropriately large.

Let $V = \{z \mid |z| < R\}$ and U be the component of $f^{-1}(V)$ which contains the origin. From (50), it follows that $U \Subset V$ and $f : U \to V$ is of degree $l+1$ and thus (U, V, f) is a polynomial-like map (whose Julia set may not be connected). From the last paragraph it follows that (U, V, f) is K-qc conjugate to some polynomial of degree $l+1$. This proves the Claim.

Now let $M_1 = 1$. By successively applying the Claim we get

$$1 = M_1 < M_2 < \cdots < M_{d-1}.$$

Let $M = M_{d-1}$. Suppose $1 = |c_{d-2}| \le \cdots \le |c_1|$.

In the first case, $|c_i| \le M$ for all $1 \le i \le d-2$. The first assertion obviously holds in this case. Note that the absolute value of the leading coefficient of f is $1/|c_1 \cdot \cdots \cdot |c_{d-1}| \ge 1/M^{d-1}$, and that all the other coefficients of f are bounded above by some constant $K(d, M)$ depending on d and M. This implies the existence of an $R(d, M)$ such that f is dominated by the leading term for all $|z| > R(d, M)$. In particular, this implies that the diameter of the Siegel disk is not greater than $R(d, M)$. Since M depends only on d, the second assertion of the lemma follows.

In the second case, there is some c_i such that $|c_i| > M$. Let $1 \le l \le d-2$ be the least integer such that $|c_{d-l-1}| > M_{l+1}$. Then we have (49). The first assertion follows from the Claim. For the second assertion, let us go back to the proof of the Claim. We see the Siegel disk centered at the origin is contained in $V = \{z \,|\, |z| < R\}$. Since $R < M_{l+1} \le M$, it follows that the diameter of the Siegel disk is not greater than $2M$. Since M depends only on d, the second assertion follows.

The proof of Lemma 7.1 is completed. $\square$

Let $0 < \alpha < 1$ be a bounded type irrational number and $d \ge 2$ be an integer. Let $\mathcal{P}_\alpha^d$ denote the class of all the polynomial maps f such that

$$f(z) = e^{2\pi i\alpha} z + \alpha_2 z + \cdots + \alpha_d z^d$$

with $\alpha_d \ne 0$ and $f'(1) = 0$. Let Δ denote the unit disk and D denote the Siegel disk of f centered at the origin.

Lemma 7.2 ([18], Shishikura). *There exists a $K = K(\alpha, d) > 1$ depending only on d and α such that for any polynomial map $f \in \mathcal{P}_\alpha^d$, if $\phi : \Delta \to D$ is the holomorphic isomorphism such that $\phi^{-1} \circ f \circ \phi(z) = e^{2\pi i\alpha} z$ for $z \in \Delta$, then ϕ can be extended to a K-qc homeomorphism of the plane. In particular, the boundary of the Siegel disk of f centered at the origin is a K-quasicircle.*

For a proof, see [18] or [23].

Lemma 7.3. *The boundaries of the Siegel disks of*

$$f \in \bigcup_{2 \le l \le d} \mathcal{P}_\alpha^l$$

at the origin moves continuously with respect to the Hausdorff metric on the spaces of non-empty compact sets of the plane and the topology of $\bigcup_{2\le l\le d} \mathcal{P}_\alpha^l$ is given by open-compact topology, that is, $f_n \to f$ with respect to this topology means that f_n uniformly converges to f in any compact set of the plane.

Proof. Let $f \in \bigcup_{2\le l\le d} \mathcal{P}_\alpha^l$. Assume that $f_n \to f$. Let D and D_n be respectively the Siegel disks of f and f_n which are centered at the origin. It suffices to prove that ∂D_g is close to ∂D_f with respect to the Hausdorff metric for all n large enough. It is known that both of them contains critical points.

We may assume that $D \neq D_n$ since otherwise there is nothing to prove. Since both D and D_n contains the origin as an interior point there is a point $w \in (D \cap \partial D_n) \cup (D_n \cap \partial D)$. Without loss of generality, let us assume that $w \in D \cap \partial Dn$. Let $\Gamma_w \subset D$ be the f-invariant curve containing w. Let $\mathcal{O}_f(w) = \{f^k(w)\}_{k \geq 0}$ and $\mathcal{O}_g(w) = \{g^k(w)\}_{k \geq 0}$. Then $\mathcal{O}_f(w)$ and $\mathcal{O}_g(w)$ are dense in Γ_w and ∂D_n respectively. For any integer $m \geq 1$, the two finite orbit segments

$$\{f^k(w), 0 \leq k \leq m\}; \quad \text{and} \quad \{f^k(w), 0 \leq k \leq m\}$$

can be arbitrarily close to each other provided that n is large enough. By Lemma 7.2 there exist two $K(\alpha, d)$-qc homeomorphisms of the plane which fix 0 and ∞, say ϕ and ψ such that $\phi^{-1} \circ f \circ \phi(z) = e^{2\pi i \alpha} z$ and $\psi^{-1} \circ g \circ \psi(z) = e^{2\pi i \alpha} z$ for all $z \in \overline{\Delta}$. $\phi(\mathbb{T}_r) = \Gamma_w$ and $\psi(\mathbb{T}) = \partial D_g$. Suppose $w = \phi(z_0)$ for some z_0 with $0 < |z_0| < 1$ and $w = \psi(\zeta_0)$ for some $\zeta_0 \in \mathbb{T}$. Then Each component of $\Gamma_w \setminus \{f^k(w), 0 \leq k \leq m\}$ is the ϕ-image of a component of

$$\{z | \, |z| = |z_0|\} \setminus \{e^{2k\pi i \alpha} \cdot z_0 \mid 0 \leq k \leq m\}$$

and each component of $\partial D_n \setminus \{f_n^k(w), 0 \leq k \leq m\}$ is the ψ-image of a component of

$$\mathbb{T} \setminus \{e^{2k\pi i \alpha} \cdot \zeta_0 \mid 0 \leq k \leq m\}.$$

By the second assertion of Lemma 7.1 the Siegel disks of f and f_n are contained in some compact set of the plane. Since ϕ and ψ fix 0 and ∞ and are K-qc homeomorphisms of the plane for some K depending only on d and α, there exist $C > 1$ and $0 < \eta < 1$ which are independent of n such that for any z_1, z_2 with $|z_1|, |z_2| \leq 1$ we have

$$|\phi(z_1) - \phi(z_2)| < C \cdot |z_1 - z_2|^\eta \text{ and } |\psi(z_1) - \psi(z_2)| < C \cdot |z_1 - z_2|^\eta.$$

Since the components of

$$\{z | \, |z| = |z_0|\} \setminus \{e^{2k\pi i \alpha} \cdot z_0 \mid 0 \leq k \leq m\}$$

and

$$\mathbb{T} \setminus \{e^{2k\pi i \alpha} \cdot \zeta_0 \mid 0 \leq k \leq m\}$$

can be arbitrarily small provided that m is large enough, it follows that Γ_w and ∂D_n can be arbitrarily to each other provided that n is large enough.

Now we claim that Γ_w is close to ∂D also provided that n is large enough. This is because if not, then by Lemma 7.2 it follows that the modulus of the annulus bounded by ∂D and Γ_w has a positive lower bound. But since ∂D_n is close to Γ_w and contains a critical point of f_n, it follows that there is a critical point of f_n contained in D and is bounded away from ∂D. But since $f_n \to f$ uniformly in D, there would be a critical point of f contained in D. This is impossible. This completes the proof of Lemma 7.3. $\square$

Lemma 7.4. *Suppose $f_n \in \bigcup_{2 \leq l \leq d} \mathcal{P}_\alpha^l$ is a sequence such that for each f_n, the boundary of the Siegel disk centered at the origin passes through the critical point 1. Then f_n has a subsequence f_{n_k} which converges to some $f \in \bigcup_{2 \leq l \leq d} \mathcal{P}_\alpha^l$. Moreover, the Siegel disk of the limit polynomial map g centered at the origin also passes through the critical point 1, and for any $k > m \geq 0$, we have*

$$\lim_{n \to \infty} \sigma_{k,m}(f_n) = \sigma_{k,m}(g).$$

Proof. By the second assertion of Lemma 7.1, the critical points of all f_n are uniformly bounded away from the origin; that us, there is a uniform $L > 0$ such that for each f_n, the critical points of f_n are contained in the outside of the disk $\{z \mid |z| > L\}$. For each f_n, let us label the critical points of f_n by

$$c_1^n, \cdots, c_{d-2}^n, c_{d-1}^n = 1.$$

By taking a subsequence, we get $1 \le i_1 < \cdots < i_l \le d-2$ and $d-1-l$ such that for all $1 \le j \le l$,

$$c_{i_j}^n \to \infty \text{ and } n \to \infty,$$

and for all $k \ne i_j, 1 \le j \le l$ and $1 \le k \le d-1$, we have

$$c_k^n \to c_k^*$$

where c_k^* is some non-zero complex number. Let g denote the polynomial of degree $d-l$ which has critical points at these $c_k^*, 1 \le k \le d-1$ and $k \ne i_j, 1 \le j \le l$. It is clear that f_{n_k} converges to g uniformly in any compact set of the plane. This proves the first assertion of Lemma 7.4. Let us prove that the boundary of the Siegel disk of g centered at the origin must also contain the critical point 1. Suppose this were not true. Then the critical point 1 is bounded away from the boundary of the Siegel disk g. Since $f_n \to g$, by Lemma 7.3, the boundary of the Siegel disk of f_n centered at the origin can be arbitrarily close to the boundary of the Siegel disk of g centered at the origin. This would imply for all n large enough, the boundary of the Siegel disk of f_n centered at the origin does not pass through the critical point 1. This is a contradiction. For $k > m \ge 0$ given, $f_n^k \to g^k$ and $f_n^m \to g^m$ uniformly in any compact set of the plane. Thus $\sigma_{k,m}(f_n) \to \sigma_{k,m}(g)$ as $n \to \infty$. This implies the second assertion. The proof of Lemma 7.4 is completed. □

References

[1] L. V. Ahlfors, *Complex Analysis*, McGRAW-HILL BOOK COMPANY, INC. 1953.

[2] A. Chéritat, *Relatively compact Siegel disks with non-locally connected boundaries*, Math. Ann. , vol. 349, no. 3, pp. 529-542, 2011.

[3] T. Chen and Y. Jiang, *Canonical Thurston obstructions for sub-hyperbolic semi-rational branched coverings.* arXiv:1101.2285v2 [math.DS]

[4] G. Cui and L. Tan, *A characterization of hyperbolic rational maps.* Invent. Math., Vol. 183 (2011), No. 3, 451-516.

[5] G. David, *Solutions de l'equation de Beltrami avec* $\|\mu\|_\infty = 1$, Ann. Acad. Sci. Fenn. Ser. A I Math., 13(1988) 25-70.

[6] E. de Faria, W. De Melo, *Rigidity of critical circle mappings I* , J. Eur. Math. Soc.(JEMS), 1(1999)339-392.

[7] A. Douady, *Disques de Siegel et anneaux de Herman*, Seminar Bourbaki, Astéerisque, 152-153 (1987) 151-172.

[8] A. Douady and J. Hubbard, *A proof of Thurston's topological characterization of rational functions*, Acta Math., 171 (1993), 263-297.

[9] M. Herman, *Uniformity of the Swiatek distorsion of the compact family of Blaschke products.* Unpublished manuscript.

[10] H. Inou and M. Shishikura, *The renormalization for parabolic fixed points and their perturbation*, manuscript, 2008.

[11] A. Ya. Khinchin, *Continued fractions*, Dover, New York, 1992.

[12] M. Lyubich, *The dynamics of raitonal transforms: the topological picture*, Russian Math. Surveys 41:4 (1986) 43-117.

[13] C. McMullen, *Self-similarity of Siegel disk and Hausdorff dimension of Julia set*, Acta Mathematica, 180 (1998), 247-292.

[14] C. Petersen, *Local connectivity of some Julia sets containing a circle with an irrational rotation*, Acta Math., 177(1996)163-224.

[15] C. Petersen, *Herman-Swiatek Theorems with Applications*, London Mathematical Society Lecture Note Series. 274 (2000), 211-225.

[16] C. Petersen, *On holomorphic critical quasi-circle maps*, Ergodic Theory Dynam. Systems 24 (2004), no. 5, 1739C1751.

[17] C. Petersen, S. Zakeri, *On the Julia set of a Typical Quadratic Polynomial with a Siegel disk*, Annals of Mathematics , Vol 159 (2004), NO. 1, 1-52.

[18] M. Shishikura, *On bounded type Siegel disks of polynomial maps.* Unpublished manuscript. 2001.

[19] P. Tukia, *Compactness properties of μ-homeomorphisms.* Ann. Acad. Sci. Fenn. Ser. A I Math., Vol. 16 (1991) 47-69.

[20] J.-C. Yoccoz, Π_0*n'y a pas de contre-exemple de Denjoy analytique*, C. R. Acad. Sci. Paris, 298(1984), 141-144.

[21] J.-C. Yoccoz, *Structure des orbites des homéomorphisms analytiques posedant un point critique*, Prepint, 1989.

[22] S. Zakeri, *Dynamics of cubic Siegel polynomials*, Comm. Math. Phys., 206 (1999) 185-233.

[23] G. Zhang, *All bounded type Siegel disks of rational maps are quasi-disks.* Invent. Math. Vol. 185, NO. 2, 421-466.

[24] G. Zhang, *On the non-escaping set of* $e^{2\pi i\theta}\sin(z)$, Israel J. Math., 165 (2008), 233-252.

[25] G. Zhang, *On David type Siegels of the sine family.* arXiv:0812.0431v2.

[26] G. Zhang and Y. Jiang, *Combinatorial characterization of sub-hyperbolic rational maps.* Advances in Mathematics, 221 (2009), 1990-2018.

Department of Mathematics, Nanjing University, Nanjing, 210093, P. R. China
E-mail address: `zhanggf@hotmail.com`